

**Phospho-MSK-S360 Antibody Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP3172a****Specification**

---

**Phospho-MSK-S360 Antibody Blocking Peptide - Product Information**Primary Accession [O75582](#)**Phospho-MSK-S360 Antibody Blocking Peptide - Additional Information****Gene ID** 9252**Other Names**

Ribosomal protein S6 kinase alpha-5, S6K-alpha-5, 90 kDa ribosomal protein S6 kinase 5, Nuclear mitogen- and stress-activated protein kinase 1, RSK-like protein kinase, RSKL, RPS6KA5, MSK1

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP3172a](/product/products/AP3172a) was selected from the region of human Phospho-MSK-S360. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**Phospho-MSK-S360 Antibody Blocking Peptide - Protein Information****Name** RPS6KA5**Synonyms** MSK1**Function**

Serine/threonine-protein kinase that is required for the mitogen or stress-induced phosphorylation of the transcription factors CREB1 and ATF1 and for the regulation of the transcription factors RELA, STAT3 and ETV1/ER81, and that contributes to gene activation by histone phosphorylation and functions in the regulation of inflammatory genes (PubMed: [11909979](http://www.uniprot.org/citations/11909979), PubMed: [12569367](http://www.uniprot.org/citations/12569367), PubMed: [12763138](http://www.uniprot.org/citations/12763138), PubMed: [9687510](http://www.uniprot.org/citations/9687510), PubMed: [18511904](http://www.uniprot.org/citations/18511904)).

[9873047](http://www.uniprot.org/citations/9873047)). Phosphorylates CREB1 and ATF1 in response to mitogenic or stress stimuli such as UV-C irradiation, epidermal growth factor (EGF) and anisomycin (PubMed:[11909979](http://www.uniprot.org/citations/11909979)), PubMed:[9873047](http://www.uniprot.org/citations/9873047)). Plays an essential role in the control of RELA transcriptional activity in response to TNF and upon glucocorticoid, associates in the cytoplasm with the glucocorticoid receptor NR3C1 and contributes to RELA inhibition and repression of inflammatory gene expression (PubMed:[12628924](http://www.uniprot.org/citations/12628924)), PubMed:[18511904](http://www.uniprot.org/citations/18511904)). In skeletal myoblasts is required for phosphorylation of RELA at 'Ser-276' during oxidative stress (PubMed:[12628924](http://www.uniprot.org/citations/12628924)). In erythropoietin-stimulated cells, is necessary for the 'Ser-727' phosphorylation of STAT3 and regulation of its transcriptional potential (PubMed:[12763138](http://www.uniprot.org/citations/12763138)). Phosphorylates ETV1/ER81 at 'Ser-191' and 'Ser-216', and thereby regulates its ability to stimulate transcription, which may be important during development and breast tumor formation (PubMed:[12569367](http://www.uniprot.org/citations/12569367)). Directly represses transcription via phosphorylation of 'Ser-1' of histone H2A (PubMed:[15010469](http://www.uniprot.org/citations/15010469)). Phosphorylates 'Ser-10' of histone H3 in response to mitogenics, stress stimuli and EGF, which results in the transcriptional activation of several immediate early genes, including proto-oncogenes c-fos/FOS and c-jun/JUN (PubMed:[12773393](http://www.uniprot.org/citations/12773393)). May also phosphorylate 'Ser-28' of histone H3 (PubMed:[12773393](http://www.uniprot.org/citations/12773393)). Mediates the mitogen- and stress-induced phosphorylation of high mobility group protein 1 (HMG1/HMG14) (PubMed:[12773393](http://www.uniprot.org/citations/12773393)). In lipopolysaccharide-stimulated primary macrophages, acts downstream of the Toll-like receptor TLR4 to limit the production of pro-inflammatory cytokines (By similarity). Functions probably by inducing transcription of the MAP kinase phosphatase DUSP1 and the anti-inflammatory cytokine interleukin 10 (IL10), via CREB1 and ATF1 transcription factors (By similarity). Plays a role in neuronal cell death by mediating the downstream effects of excitotoxic injury (By similarity). Phosphorylates TRIM7 at 'Ser-107' in response to growth factor signaling via the MEK/ERK pathway, thereby stimulating its ubiquitin ligase activity (PubMed:[25851810](http://www.uniprot.org/citations/25851810)).

#### **Cellular Location**

Nucleus. Cytoplasm. Note=Predominantly nuclear. Exported into cytoplasm in response to glucocorticoid

#### **Tissue Location**

Widely expressed with high levels in heart, brain and placenta. Less abundant in lung, kidney and liver

### **Phospho-MSK-S360 Antibody Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

### **Phospho-MSK-S360 Antibody Blocking Peptide - Images**

### **Phospho-MSK-S360 Antibody Blocking Peptide - Background**

MSK1 is a serine/threonine kinase that may play a role in mediating the growth-factor and stress induced activation of the transcription factor CREB. This protein plays an essential role in the

control of RELA transcriptional activity in response to TNF. Activity of MSK1 requires the presence of both kinase domains.

#### **Phospho-MSK-S360 Antibody Blocking Peptide - References**

Beausoleil, S.A., et al., Proc. Natl. Acad. Sci. U.S.A. 101(33):12130-12135 (2004). Abecassis, L., et al., J. Biol. Chem. 279(29):30474-30479 (2004). Zhang, Y., et al., Carcinogenesis 25(7):1165-1175 (2004). Zhang, Y., et al., J. Biol. Chem. 279(21):21866-21872 (2004). Li, Y., et al., J. Biol. Chem. 279(2):970-979 (2004).