

Phospho-PIK3R1(Y368) Antibody Blocking peptide
Synthetic peptide
Catalog # BP3336a**Specification**

Phospho-PIK3R1(Y368) Antibody Blocking peptide - Product InformationPrimary Accession [P27986](#)**Phospho-PIK3R1(Y368) Antibody Blocking peptide - Additional Information****Gene ID** 5295**Other Names**

Phosphatidylinositol 3-kinase regulatory subunit alpha, PI3-kinase regulatory subunit alpha, PI3K regulatory subunit alpha, PtdIns-3-kinase regulatory subunit alpha, Phosphatidylinositol 3-kinase 85 kDa regulatory subunit alpha, PI3-kinase subunit p85-alpha, PtdIns-3-kinase regulatory subunit p85-alpha, PIK3R1, GRB1

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP3336a](/product/products/AP3336a) was selected from the region of human Phospho-PIK3R1-Y368. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

Phospho-PIK3R1(Y368) Antibody Blocking peptide - Protein Information**Name** PIK3R1**Synonyms** GRB1**Function**

Binds to activated (phosphorylated) protein-Tyr kinases, through its SH2 domain, and acts as an adapter, mediating the association of the p110 catalytic unit to the plasma membrane. Necessary for the insulin-stimulated increase in glucose uptake and glycogen synthesis in insulin-sensitive tissues. Plays an important role in signaling in response to FGFR1, FGFR2, FGFR3, FGFR4, KITLG/SCF, KIT, PDGFRA and PDGFRB. Likewise, plays a role in ITGB2 signaling (PubMed: [17626883](http://www.uniprot.org/citations/17626883), PubMed: [19805105](http://www.uniprot.org/citations/19805105), PubMed: [19805105](http://www.uniprot.org/citations/19805105)).

[7518429](http://www.uniprot.org/citations/7518429)). Modulates the cellular response to ER stress by promoting nuclear translocation of XBP1 isoform 2 in a ER stress- and/or insulin-dependent manner during metabolic overloading in the liver and hence plays a role in glucose tolerance improvement (PubMed: [20348923](http://www.uniprot.org/citations/20348923)).

Tissue Location

Isoform 2 is expressed in skeletal muscle and brain, and at lower levels in kidney and cardiac muscle. Isoform 2 and isoform 4 are present in skeletal muscle (at protein level)

Phospho-PIK3R1(Y368) Antibody Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

Phospho-PIK3R1(Y368) Antibody Blocking peptide - Images

Phospho-PIK3R1(Y368) Antibody Blocking peptide - Background

Phosphatidylinositol 3-kinase phosphorylates the inositol ring of phosphatidylinositol at the 3-prime position. The enzyme comprises a 110 kD catalytic subunit and a regulatory subunit of either 85, 55, or 50 kD. This antibody is directed against an epitope in the 85 kD regulatory subunit. Phosphatidylinositol 3-kinase plays an important role in the metabolic actions of insulin, and a mutation in the gene has been associated with insulin resistance.

Phospho-PIK3R1(Y368) Antibody Blocking peptide - References

Kobayashi, H., et al., J. Biol. Chem. 279(8):6371-6379 (2004). Liu, H., et al., J. Cell Biol. 164(4):603-612 (2004). Sun, M., et al., J. Biol. Chem. 278(44):42992-43000 (2003). Khan, N.A., et al., J. Neurovirol. 9(6):584-593 (2003). Lee, H.Y., et al., J. Biol. Chem. 278(26):23630-23638 (2003).