

Phospho-HSPB1(S83) Antibody Blocking peptide Synthetic peptide Catalog # BP3431a

Specification

Phospho-HSPB1(S83) Antibody Blocking peptide - Product Information

Primary Accession

<u>P04792</u>

Phospho-HSPB1(S83) Antibody Blocking peptide - Additional Information

Gene ID 3315

Other Names

Heat shock protein beta-1, HspB1, 28 kDa heat shock protein, Estrogen-regulated 24 kDa protein, Heat shock 27 kDa protein, HSP 27, Stress-responsive protein 27, SRP27, HSPB1, HSP27, HSP28

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP3431a was selected from the region of human Phospho-HSPB1-pS83. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions This product is for research use only. Not for use in diagnostic or therapeutic procedures.

Phospho-HSPB1(S83) Antibody Blocking peptide - Protein Information

Name HSPB1

Synonyms HSP27, HSP28

Function

Small heat shock protein which functions as a molecular chaperone probably maintaining denatured proteins in a folding- competent state (PubMed:10383393, PubMed:20178975). Plays a role
in stress resistance and actin organization (PubMed:<a/pre>

href="http://www.uniprot.org/citations/19166925" target="_blank">19166925). Through its molecular chaperone activity may regulate numerous biological processes including the phosphorylation and the axonal transport of neurofilament proteins (PubMed:23728742).



Cellular Location

Cytoplasm. Nucleus Cytoplasm, cytoskeleton, spindle Note=Cytoplasmic in interphase cells. Colocalizes with mitotic spindles in mitotic cells. Translocates to the nucleus during heat shock and resides in sub-nuclear structures known as SC35 speckles or nuclear splicing speckles.

Tissue Location

Detected in all tissues tested: skeletal muscle, heart, aorta, large intestine, small intestine, stomach, esophagus, bladder, adrenal gland, thyroid, pancreas, testis, adipose tissue, kidney, liver, spleen, cerebral cortex, blood serum and cerebrospinal fluid. Highest levels are found in the heart and in tissues composed of striated and smooth muscle.

Phospho-HSPB1(S83) Antibody Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

<u>Blocking Peptides</u>

Phospho-HSPB1(S83) Antibody Blocking peptide - Images

Phospho-HSPB1(S83) Antibody Blocking peptide - Background

In response to adverse changes in their environment, cells from many organisms increase the expression of a class of proteins referred to as heat shock or stress proteins. HSBP1 exhibits rapid increased phosphorylation in response to various mitogens, tumor promoters (e.g. phorbol esters) and calcium ionophores, and high levels are associated with carcinoma of the breast and with endometrial adenocarcinomas. Heat shock of HeLa cell cultures, or treatment with arsenite, phorbol ester, or tumor necrosis factor, causes a rapid phosphorylation of preexisting HSBP1, with Ser82 as the major site and Ser78 the minor site of phosphorylation. HSBP1 may exert phosphorylation-activated functions linked with growth signaling pathways in unstressed cells. A homeostatic function at this level could protect cells from adverse effects of signal transduction systems which may be activated inappropriately during stress.

Phospho-HSPB1(S83) Antibody Blocking peptide - References

Wano, C., et al., Exp. Cell Res. 298(2):584-592 (2004).Evgrafov, O.V., et al., Nat. Genet. 36(6):602-606 (2004).Song, H., et al., Biochem. Biophys. Res. Commun. 314(1):143-150 (2004).Chauhan, D., et al., Blood 102(9):3379-3386 (2003).Van Why, S.K., et al., J. Am. Soc. Nephrol. 14(1):98-106 (2003).