

Phospho-ENT1(Slc29a1)(S254) Antibody Blocking peptide
Synthetic peptide
Catalog # BP3615a**Specification**

Phospho-ENT1(Slc29a1)(S254) Antibody Blocking peptide - Product InformationPrimary Accession [Q99808](#)**Phospho-ENT1(Slc29a1)(S254) Antibody Blocking peptide - Additional Information****Gene ID** 2030**Other Names**

Equilibrative nucleoside transporter 1, Equilibrative nitrobenzylmercaptapurine riboside-sensitive nucleoside transporter, Equilibrative NBMPR-sensitive nucleoside transporter, Nucleoside transporter, es-type, Solute carrier family 29 member 1, SLC29A1, ENT1

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP3615a](/products/AP3615a) was selected from the Slc29a1 region of human Phospho-ENT1(Slc29a1). A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

Phospho-ENT1(Slc29a1)(S254) Antibody Blocking peptide - Protein Information**Name** SLC29A1 ([HGNC:11003](#))**Synonyms** ENT1**Function**

Uniporter involved in the facilitative transport of nucleosides and nucleobases, and contributes to maintaining their cellular homeostasis (PubMed:[8986748](http://www.uniprot.org/citations/8986748), PubMed:[10755314](http://www.uniprot.org/citations/10755314), PubMed:[12527552](http://www.uniprot.org/citations/12527552), PubMed:[10722669](http://www.uniprot.org/citations/10722669), PubMed:[21795683](http://www.uniprot.org/citations/21795683), PubMed:[35790189](http://www.uniprot.org/citations/35790189), PubMed:[35790189](http://www.uniprot.org/citations/35790189)

href="http://www.uniprot.org/citations/27995448" target="_blank">27995448, PubMed:17379602, PubMed:14759222, PubMed:15037197, PubMed:26406980). Functions as a Na(+)-independent transporter (PubMed:8986748). Involved in the transport of nucleosides such as adenosine, guanosine, inosine, uridine, thymidine and cytidine (PubMed:8986748, PubMed:10755314, PubMed:12527552, PubMed:10722669, PubMed:17379602, PubMed:14759222, PubMed:15037197, PubMed:26406980). Also transports purine nucleobases (hypoxanthine, adenine, guanine) and pyrimidine nucleobases (thymine, uracil) (PubMed:21795683, PubMed:27995448). Mediates basolateral nucleoside uptake into Sertoli cells, thereby regulating the transport of nucleosides in testis across the blood-testis barrier (By similarity). Regulates inosine levels in brown adipocytes tissues (BAT) and extracellular inosine levels, which controls BAT-dependent energy expenditure (PubMed:35790189).

Cellular Location

Basolateral cell membrane; Multi-pass membrane protein. Apical cell membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein. Note=Localized to the basolateral membrane of Sertoli cells (PubMed:23639800). Localized to the cell membrane of erythrocytes (PubMed:23219802, PubMed:11584005).

Tissue Location

Expressed in testis at the blood-testis barrier (at protein level) (PubMed:23639800). Detected in erythrocytes (at protein level) (PubMed:23219802, PubMed:11584005). Expressed at relatively high levels in cerebral cortex, particularly the frontal and parietal lobes, and the thalamus and basal ganglia (at protein level) (PubMed:11311901). In the midbrain expressed at moderate levels, whereas in the other areas of the brainstem, namely medulla and pons, cerebellum and the hippocampus expressed at lower amounts when compared to the other brain regions (at protein level) (PubMed:11311901). Expressed in Langerhans cells and lymphocytes in the pancreas (at protein level) (PubMed:15501974). Expressed in kidney, in polarized renal epithelial cells (PubMed:12527552). Expressed in adipose tissues (PubMed:35790189). Expressed in placenta (PubMed:8986748). Expressed in small intestine (PubMed:10755314).

Phospho-ENT1(Slc29a1)(S254) Antibody Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

Phospho-ENT1(Slc29a1)(S254) Antibody Blocking peptide - Images

Phospho-ENT1(Slc29a1)(S254) Antibody Blocking peptide - Background

ENT1 is a member of the equilibrative nucleoside transporter family. The protein is categorized as an equilibrative (as opposed to concentrative) transporter that is sensitive to inhibition by nitrobenzylthioinosine (NBMPR). Nucleoside transporters are required for nucleotide synthesis in cells that lack de novo nucleoside synthesis pathways, and are also necessary for the uptake of

cytotoxic nucleosides used for cancer and viral chemotherapies.

Phospho-ENT1(Slc29a1)(S254) Antibody Blocking peptide - References

Dephoure N, et al. (2008) Proc Natl Acad Sci U S A 105, 10762-7
Bone DB, Robillard KR, Stolk M, Hammond JR (2007) Mol Membr Biol 24, 294-303