

Phospho-H3(S10) Antibody Blocking peptide Synthetic peptide

Catalog # BP3667a

Specification

Phospho-H3(S10) Antibody Blocking peptide - Product Information

Primary Accession

<u>P84243</u>

Phospho-H3(S10) Antibody Blocking peptide - Additional Information

Gene ID 3020;3021

Other Names Histone H33, H3F3A, H33A, H3F3

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP3667a was selected from the region of human Phospho-H3-S10. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions This product is for research use only. Not for use in diagnostic or therapeutic procedures.

Phospho-H3(S10) Antibody Blocking peptide - Protein Information

Name H3-3A (<u>HGNC:4764</u>)

Synonyms H3.3A, H3F3, H3F3A

Function

Variant histone H3 which replaces conventional H3 in a wide range of nucleosomes in active genes. Constitutes the predominant form of histone H3 in non-dividing cells and is incorporated into chromatin independently of DNA synthesis. Deposited at sites of nucleosomal displacement throughout transcribed genes, suggesting that it represents an epigenetic imprint of transcriptionally active chromatin. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.



Cellular Location Nucleus. Chromosome

Phospho-H3(S10) Antibody Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

Blocking Peptides

Phospho-H3(S10) Antibody Blocking peptide - Images

Phospho-H3(S10) Antibody Blocking peptide - Background

H3 is basic nuclear protein that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A,H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures.

Phospho-H3(S10) Antibody Blocking peptide - References

Teng,H., et.al., J. Biol. Chem. 284 (39), 26368-26376 (2009)Garcia,B.A., et.al., J. Biol. Chem. 282 (10), 7641-7655 (2007)