

# Phospho-Endophilin(Y80) Antibody Blocking peptide

Synthetic peptide Catalog # BP3682a

# **Specification**

## Phospho-Endophilin(Y80) Antibody Blocking peptide - Product Information

**Primary Accession** 

**09Y371** 

# Phospho-Endophilin(Y80) Antibody Blocking peptide - Additional Information

**Gene ID 51100** 

#### **Other Names**

Endophilin-B1, Bax-interacting factor 1, Bif-1, SH3 domain-containing GRB2-like protein B1, SH3GLB1, KIAA0491

# **Target/Specificity**

The synthetic peptide sequence used to generate the antibody <a href=/products/AP3682a>AP3682a</a> was selected from the region of human Phospho-Endophilin-pY80. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

# Phospho-Endophilin(Y80) Antibody Blocking peptide - Protein Information

Name SH3GLB1

Synonyms KIAA0491

## **Function**

May be required for normal outer mitochondrial membrane dynamics (PubMed:<a href="http://www.uniprot.org/citations/15452144" target="\_blank">15452144</a>). Required for coatomer-mediated retrograde transport in certain cells (By similarity). May recruit other proteins to membranes with high curvature. May promote membrane fusion (PubMed:<a href="http://www.uniprot.org/citations/11604418" target="\_blank">11604418</a>). Involved in activation of caspase-dependent apoptosis by promoting BAX/BAK1 activation (PubMed:<a href="http://www.uniprot.org/citations/16227588" target="\_blank">16227588</a>). Isoform 1 acts proapoptotic in fibroblasts (By similarity). Involved in caspase- independent apoptosis during nutrition starvation and involved in the regulation of autophagy. Activates lipid kinase activity of



PIK3C3 during autophagy probably by associating with the PI3K complex II (PI3KC3-C2) (PubMed:<a href="http://www.uniprot.org/citations/17891140" target="\_blank">17891140</a>). Associated with PI3KC3-C2 during autophagy may regulate the trafficking of ATG9A from the Golgi complex to the peripheral cytoplasm for the formation of autophagosomes by inducing Golgi membrane tubulation and fragmentation (PubMed:<a

href="http://www.uniprot.org/citations/21068542" target="\_blank">21068542</a>). Involved in regulation of degradative endocytic trafficking and cytokinesis, probably in the context of PI3KC3-C2 (PubMed:<a href="http://www.uniprot.org/citations/20643123"

target="\_blank">20643123</a>). Isoform 2 acts antiapoptotic in neuronal cells; involved in maintenance of mitochondrial morphology and promotes neuronal viability (By similarity).

## **Cellular Location**

Cytoplasm. Golgi apparatus membrane; Peripheral membrane protein. Mitochondrion outer membrane; Peripheral membrane protein. Cytoplasmic vesicle, autophagosome membrane. Midbody. Note=Association with the Golgi apparatus depends on the cell type (By similarity). Following starvation colocalizes with ATG5 and LC3 autophagy-related protein(s)on autophagosomal membranes (PubMed:17891140). {ECO:0000250, ECO:0000269|PubMed:17891140}

#### **Tissue Location**

Highly expressed in heart, skeletal muscle, kidney and placenta. Detected at lower levels in brain, colon, thymus, spleen, liver, small intestine, lung and peripheral blood leukocytes

## Phospho-Endophilin(Y80) Antibody Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

# • Blocking Peptides

Phospho-Endophilin(Y80) Antibody Blocking peptide - Images

#### Phospho-Endophilin(Y80) Antibody Blocking peptide - Background

Endophilin may be required for normal outer mitochondrial membrane dynamics. It is required for coatomer-mediated retrograde transport in certain cells. It may recruit other proteins to membranes with high curvature and may promote membrane fusion.

# Phospho-Endophilin(Y80) Antibody Blocking peptide - References

Maiuri, M.C., et.al., Cell Death Differ. 16 (1), 87-93 (2009) Yamaguchi, H., et.al., J. Biol. Chem. 283 (27), 19112-19118 (2008) Takahashi, Y., et.al, Nat. Cell Biol. 9 (10), 1142-1151 (2007)