

Phospho-LIMK1(Thr508)) Antibody Blocking peptide
Synthetic peptide
Catalog # BP3745a**Specification**

Phospho-LIMK1(Thr508)) Antibody Blocking peptide - Product InformationPrimary Accession [P53667](#)**Phospho-LIMK1(Thr508)) Antibody Blocking peptide - Additional Information****Gene ID** 3984**Other Names**

LIM domain kinase 1, LIMK-1, LIMK1, LIMK

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

Phospho-LIMK1(Thr508)) Antibody Blocking peptide - Protein Information**Name** LIMK1**Synonyms** LIMK**Function**

Serine/threonine-protein kinase that plays an essential role in the regulation of actin filament dynamics. Acts downstream of several Rho family GTPase signal transduction pathways (PubMed:10436159, PubMed:11832213, PubMed:12807904, PubMed:15660133, PubMed:16230460, PubMed:18028908, PubMed:22328514, PubMed:23633677). Activated by upstream kinases including ROCK1, PAK1 and PAK4, which phosphorylate LIMK1 on a threonine residue located in its activation loop (PubMed:10436159). LIMK1 subsequently phosphorylates and inactivates the actin binding/depolymerizing factors cofilin-1/CFL1, cofilin-2/CFL2 and destrin/DSTN, thereby preventing the cleavage of filamentous actin (F-actin), and stabilizing the actin cytoskeleton (PubMed:10436159).

href="http://www.uniprot.org/citations/11832213" target="_blank">11832213, PubMed:15660133, PubMed:16230460, PubMed:23633677). In this way LIMK1 regulates several actin-dependent biological processes including cell motility, cell cycle progression, and differentiation (PubMed:11832213, PubMed:15660133, PubMed:16230460, PubMed:23633677). Phosphorylates TPPP on serine residues, thereby promoting microtubule disassembly (PubMed:18028908). Stimulates axonal outgrowth and may be involved in brain development (PubMed:18028908).

Cellular Location

Cytoplasm. Nucleus. Cytoplasm, cytoskeleton. Cell projection, lamellipodium {ECO:0000250|UniProtKB:P53668} Note=Predominantly found in the cytoplasm. Localizes in the lamellipodium in a CDC42BPA, CDC42BPB and FAM89B/LRAP25-dependent manner. {ECO:0000250|UniProtKB:P53668}

Tissue Location

Highest expression in both adult and fetal nervous system. Detected ubiquitously throughout the different regions of adult brain, with highest levels in the cerebral cortex. Expressed to a lesser extent in heart and skeletal muscle

Phospho-LIMK1(Thr508)) Antibody Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

Phospho-LIMK1(Thr508)) Antibody Blocking peptide - Images

Phospho-LIMK1(Thr508)) Antibody Blocking peptide - Background

There are approximately 40 known eukaryotic LIM proteins,so named for the LIM domains they contain. LIM domains are highlyconserved cysteine-rich structures containing 2 zinc fingers.Although zinc fingers usually function by binding to DNA or RNA,the LIM motif probably mediates protein-protein interactions. LIMkinase-1 and LIM kinase-2 belong to a small subfamily with a uniquecombination of 2 N-terminal LIM motifs and a C-terminal proteinkinase domain. LIMK1 is likely to be a component of anintracellular signaling pathway and may be involved in braindevelopment. LIMK1 hemizygoty is implicated in the impairedvisuospatial constructive cognition of Williams syndrome. [providedby RefSeq].

Phospho-LIMK1(Thr508)) Antibody Blocking peptide - References

There are approximately 40 known eukaryotic LIM proteins,so named for the LIM domains they contain. LIM domains are highlyconserved cysteine-rich structures containing 2 zinc fingers.Although zinc fingers usually function by binding to DNA or RNA,the LIM motif probably mediates protein-protein interactions. LIMkinase-1 and LIM kinase-2 belong to a small subfamily with a uniquecombination of 2 N-terminal LIM motifs and a C-terminal proteinkinase domain. LIMK1 is likely to be a component of anintracellular signaling pathway and may be involved in braindevelopment. LIMK1 hemizygoty is implicated in the impairedvisuospatial constructive cognition of Williams syndrome. [providedby RefSeq].