

Phospho-IKKB(S692) Blocking Peptide
Synthetic peptide
Catalog # BP3782f**Specification**

Phospho-IKKB(S692) Blocking Peptide - Product Information

Primary Accession [O14920](#)
Other Accession [NP_001177649.1](#)

Phospho-IKKB(S692) Blocking Peptide - Additional Information

Gene ID 3551

Other Names

Inhibitor of nuclear factor kappa-B kinase subunit beta, I-kappa-B-kinase beta, IKK-B, IKK-beta, IKBKB, I-kappa-B kinase 2, IKK2, Nuclear factor NF-kappa-B inhibitor kinase beta, NFKBIKB, IKBKB, IKKB

Target/Specificity

The synthetic peptide sequence is selected from aa 685-699 of HUMAN IKBKB

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

Phospho-IKKB(S692) Blocking Peptide - Protein Information

Name IKBKB

Synonyms IKKB

Function

Serine kinase that plays an essential role in the NF-kappa-B signaling pathway which is activated by multiple stimuli such as inflammatory cytokines, bacterial or viral products, DNA damages or other cellular stresses (PubMed: [20434986](http://www.uniprot.org/citations/20434986), PubMed: [20797629](http://www.uniprot.org/citations/20797629), PubMed: [21138416](http://www.uniprot.org/citations/21138416), PubMed: [9346484](http://www.uniprot.org/citations/9346484), PubMed: [30337470](http://www.uniprot.org/citations/30337470)). Acts as a part of the canonical IKK complex in the conventional pathway of NF-kappa-B activation (PubMed: [9346484](http://www.uniprot.org/citations/9346484), PubMed: [30337470](http://www.uniprot.org/citations/30337470)). Phosphorylates inhibitors of NF-kappa-B on 2 critical serine

residues (PubMed:9346484, PubMed:20434986, PubMed:20797629, PubMed:21138416). These modifications allow polyubiquitination of the inhibitors and subsequent degradation by the proteasome (PubMed:9346484, PubMed:20434986, PubMed:20797629, PubMed:21138416). In turn, free NF-kappa-B is translocated into the nucleus and activates the transcription of hundreds of genes involved in immune response, growth control, or protection against apoptosis (PubMed:9346484, PubMed:20434986, PubMed:20797629, PubMed:21138416). In addition to the NF-kappa-B inhibitors, phosphorylates several other components of the signaling pathway including NEMO/IKBKG, NF-kappa-B subunits RELA and NFkB1, as well as IKK-related kinases TBK1 and IKKBE (PubMed:11297557, PubMed:14673179, PubMed:20410276, PubMed:21138416). IKK-related kinase phosphorylations may prevent the overproduction of inflammatory mediators since they exert a negative regulation on canonical IKKs (PubMed:11297557, PubMed:20410276, PubMed:21138416). Phosphorylates FOXO3, mediating the TNF-dependent inactivation of this pro-apoptotic transcription factor (PubMed:15084260). Also phosphorylates other substrates including NAA10, NCOA3, BCL10 and IRS1 (PubMed:19716809, PubMed:17213322). Phosphorylates RIPK1 at 'Ser-25' which represses its kinase activity and consequently prevents TNF- mediated RIPK1-dependent cell death (By similarity). Phosphorylates the C-terminus of IRF5, stimulating IRF5 homodimerization and translocation into the nucleus (PubMed:25326418).

Cellular Location

Cytoplasm. Nucleus. Membrane raft. Note=Colocalized with DPP4 in membrane rafts.

Tissue Location

Highly expressed in heart, placenta, skeletal muscle, kidney, pancreas, spleen, thymus, prostate, testis and peripheral blood

Phospho-IKKB(S692) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

Phospho-IKKB(S692) Blocking Peptide - Images

Phospho-IKKB(S692) Blocking Peptide - Background

NFKB1 (MIM 164011) or NFKB2 (MIM 164012) is bound to REL (MIM 164910), RELA (MIM 164014), or RELB (MIM 604758) to form the

NFKB complex. The NFKB complex is inhibited by I-kappa-B proteins (NFKBIA, MIM 164008, or NFKBIB, MIM 604495), which inactivate NF-kappa-B by trapping it in the cytoplasm. Phosphorylation of serine residues on the I-kappa-B proteins by kinases (IKBKA, MIM 600664, or IKBKB) marks them for destruction via the ubiquitination pathway, thereby allowing activation of the NF-kappa-B complex. Activated NFKB complex translocates into the nucleus and binds DNA at kappa-B-binding motifs such as 5-prime GGGRNNYYCC 3-prime or 5-prime HGGARNYYCC 3-prime (where H is A, C, or T; R is an A or G purine; and Y is a C or T pyrimidine).

Phospho-IKKB(S692) Blocking Peptide - References

Yatherajam, G., et al. J. Immunol. 185(5):2665-2669(2010)
Kenneth, N.S., et al. EMBO J. 29(17):2966-2978(2010)
Zhao, M., et al. J. Biol. Chem. 285(32):24372-24380(2010)
Niida, M., et al. Mol. Immunol. 47(14):2378-2387(2010)
Schuurhof, A., et al. Pediatr. Pulmonol. 45(6):608-613(2010)