

**CLEC2B Antibody (Center) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP4770c****Specification**

---

**CLEC2B Antibody (Center) Blocking Peptide - Product Information**

Primary Accession [Q92478](#)

**CLEC2B Antibody (Center) Blocking Peptide - Additional Information**

**Gene ID** 9976

**Other Names**

C-type lectin domain family 2 member B, Activation-induced C-type lectin, C-type lectin superfamily member 2, IFN-alpha-2b-inducing-related protein 1, CLEC2B, AICL, CLECSF2, IFNRG1

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**CLEC2B Antibody (Center) Blocking Peptide - Protein Information**

**Name** CLEC2B

**Synonyms** AICL, CLECSF2, IFNRG1

**Cellular Location**

Membrane; Single-pass type II membrane protein

**Tissue Location**

Expressed preferentially in lymphoid tissues, and in most hematopoietic cell types

**CLEC2B Antibody (Center) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**CLEC2B Antibody (Center) Blocking Peptide - Images****CLEC2B Antibody (Center) Blocking Peptide - Background**

CLEC2B encodes a member of the C-type lectin/C-type lectin-like domain (CTL/CTLD) superfamily. Members of this family share a common protein fold and have diverse functions, such as cell adhesion, cell-cell signalling, glycoprotein turnover, and roles in inflammation and immune response. The encoded type 2 transmembrane protein may function as a cell activation antigen.

#### **CLEC2B Antibody (Center) Blocking Peptide - References**

Kuttruff, S., et al. Blood 113(2):358-369(2009)Ebner, S., et al. Proteins 53(1):44-55(2003)Renedo, M., et al. Genomics 65(2):129-136(2000)