

ATP2A1 Antibody (C-term) Blocking Peptide

Synthetic peptide Catalog # BP4905b

Specification

ATP2A1 Antibody (C-term) Blocking Peptide - Product Information

Primary Accession

014983

ATP2A1 Antibody (C-term) Blocking Peptide - Additional Information

Gene ID 487

Other Names

Sarcoplasmic/endoplasmic reticulum calcium ATPase 1, SERCA1, SR Ca(2+)-ATPase 1, Calcium pump 1, Calcium-transporting ATPase sarcoplasmic reticulum type, fast twitch skeletal muscle isoform, Endoplasmic reticulum class 1/2 Ca(2+) ATPase, ATP2A1

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

ATP2A1 Antibody (C-term) Blocking Peptide - Protein Information

Name ATP2A1 (HGNC:811)

Function

Key regulator of striated muscle performance by acting as the major Ca(2+) ATPase responsible for the reuptake of cytosolic Ca(2+) into the sarcoplasmic reticulum. Catalyzes the hydrolysis of ATP coupled with the translocation of calcium from the cytosol to the sarcoplasmic reticulum lumen (By similarity). Contributes to calcium sequestration involved in muscular excitation/contraction (PubMed:10914677).

Cellular Location

Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:P04191}; Multi-pass membrane protein {ECO:0000250|UniProtKB:P04191}. Sarcoplasmic reticulum membrane {ECO:0000250|UniProtKB:P04191}; Multi-pass membrane protein {ECO:0000250|UniProtKB:P04191}

Tissue Location

Skeletal muscle, fast twitch muscle (type II) fibers.



ATP2A1 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides

ATP2A1 Antibody (C-term) Blocking Peptide - Images

ATP2A1 Antibody (C-term) Blocking Peptide - Background

ATP2A1 encodes one of the SERCA Ca(2+)-ATPases, which are intracellular pumps located in the sarcoplasmic or endoplasmic reticula of muscle cells. This enzyme catalyzes the hydrolysis of ATP coupled with the translocation of calcium from the cytosol to the sarcoplasmic reticulum lumen, and is involved in muscular excitation and contraction.

ATP2A1 Antibody (C-term) Blocking Peptide - References

Salanova, M., et al. Histochem. Cell Biol. 132(4):383-394(2009)Thorleifsson, G., et al. Nat. Genet. 41(1):18-24(2009)Chami, M., et al. Mol. Cell 32(5):641-651(2008)