

**Tnks Blocking Peptide (C-term)**  
**Synthetic peptide**  
**Catalog # BP5358b**

### Specification

---

#### Tnks Blocking Peptide (C-term) - Product Information

Primary Accession  
Other Accession

[O95271](#)  
[Q6PFX9](#), [NP\\_003738.2](#)

#### Tnks Blocking Peptide (C-term) - Additional Information

##### Gene ID 8658

##### Other Names

Tankyrase-1, TANK1, ADP-ribosyltransferase diphtheria toxin-like 5, ARTD5, Poly [ADP-ribose] polymerase 5A, TNKS-1, TRF1-interacting ankyrin-related ADP-ribose polymerase, Tankyrase I, TNKS, PARP5A, PARPL, TIN1, TINF1, TNKS1

##### Target/Specificity

The synthetic peptide sequence is selected from aa 1305-1320 of HUMAN TNKS

##### Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

##### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

##### Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

#### Tnks Blocking Peptide (C-term) - Protein Information

##### Name TNKS ([HGNC:11941](#))

##### Function

Poly-ADP-ribosyltransferase involved in various processes such as Wnt signaling pathway, telomere length and vesicle trafficking (PubMed:<a href="http://www.uniprot.org/citations/10988299" target="\_blank">10988299</a>, PubMed:<a href="http://www.uniprot.org/citations/11739745" target="\_blank">11739745</a>, PubMed:<a href="http://www.uniprot.org/citations/16076287" target="\_blank">16076287</a>, PubMed:<a href="http://www.uniprot.org/citations/19759537" target="\_blank">19759537</a>, PubMed:<a href="http://www.uniprot.org/citations/21478859" target="\_blank">21478859</a>, PubMed:<a href="http://www.uniprot.org/citations/22864114" target="\_blank">22864114</a>, PubMed:<a href="http://www.uniprot.org/citations/23622245" target="\_blank">23622245</a>, PubMed:<a href="http://www.uniprot.org/citations/25043379" target="\_blank">25043379</a>, PubMed:<a href="http://www.uniprot.org/citations/28619731" target="\_blank">28619731</a>). Acts as an activator of the Wnt signaling pathway by mediating poly-ADP-ribosylation (PARsylation) of AXIN1

and AXIN2, 2 key components of the beta-catenin destruction complex: poly-ADP- ribosylated target proteins are recognized by RNF146, which mediates their ubiquitination and subsequent degradation (PubMed:<a href="http://www.uniprot.org/citations/19759537" target="\_blank">19759537</a>, PubMed:<a href="http://www.uniprot.org/citations/21478859" target="\_blank">21478859</a>). Also mediates PARsylation of BLZF1 and CASC3, followed by recruitment of RNF146 and subsequent ubiquitination (PubMed:<a href="http://www.uniprot.org/citations/21478859" target="\_blank">21478859</a>). Mediates PARsylation of TERF1, thereby contributing to the regulation of telomere length (PubMed:<a href="http://www.uniprot.org/citations/11739745" target="\_blank">11739745</a>). Involved in centrosome maturation during prometaphase by mediating PARsylation of HEPACAM2/MIKI (PubMed:<a href="http://www.uniprot.org/citations/22864114" target="\_blank">22864114</a>). May also regulate vesicle trafficking and modulate the subcellular distribution of SLC2A4/GLUT4-vesicles (PubMed:<a href="http://www.uniprot.org/citations/10988299" target="\_blank">10988299</a>). May be involved in spindle pole assembly through PARsylation of NUMA1 (PubMed:<a href="http://www.uniprot.org/citations/16076287" target="\_blank">16076287</a>). Stimulates 26S proteasome activity (PubMed:<a href="http://www.uniprot.org/citations/23622245" target="\_blank">23622245</a>).

### **Cellular Location**

Cytoplasm. Golgi apparatus membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Nucleus, nuclear pore complex. Chromosome, telomere. Cytoplasm, cytoskeleton, spindle pole. Note=Associated with the Golgi and with juxtanuclear SLC2A4/GLUT4-vesicles (PubMed:22864114). A minor proportion is also found at nuclear pore complexes and around the pericentriolar matrix of mitotic centromeres (PubMed:10523501). During interphase, a small fraction of TNKS is found in the nucleus, associated with TERF1 (PubMed:12768206). Localizes to spindle poles at mitosis onset via interaction with NUMA1 (PubMed:12080061)

### **Tissue Location**

Ubiquitous; highest levels in testis.

### **Tnks Blocking Peptide (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

### **Tnks Blocking Peptide (C-term) - Images**

### **Tnks Blocking Peptide (C-term) - References**

Davila, S., et al. Genes Immun. (2010) In press :  
Shen, J., et al. Cancer Epidemiol. Biomarkers Prev. 19(1):219-228(2010)  
Appierto, V., et al. Mol. Cancer Ther. 8(12):3360-3368(2009)  
Varadi, V., et al. Eur. J. Cancer 45(17):3008-3016(2009)  
Choi, J.E., et al. Cancer Epidemiol. Biomarkers Prev. 18(10):2773-2781(2009)  
Lyons, R.J., et al. J. Biol. Chem. 276(20):17172-17180(2001)  
Chi, N.W., et al. J. Biol. Chem. 275(49):38437-38444(2000)  
Smith, S., et al. J. Cell. Sci. 112 (PT 21), 3649-3656 (1999) :  
Zhu, L., et al. Genomics 57(2):320-321(1999)  
Smith, S., et al. Science 282(5393):1484-1487(1998)