

# AAA1 Antibody (C-term) Blocking peptide

Synthetic peptide Catalog # BP5508b

### **Specification**

# AAA1 Antibody (C-term) Blocking peptide - Product Information

Primary Accession Q9NS82
Other Accession NP 062823.1

# AAA1 Antibody (C-term) Blocking peptide - Additional Information

Gene ID 56301

#### **Other Names**

Asc-type amino acid transporter 1, Asc-1, Solute carrier family 7 member 10, SLC7A10, ASC1

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

### **Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

### AAA1 Antibody (C-term) Blocking peptide - Protein Information

Name SLC7A10

Synonyms ASC1

### **Function**

Associates with SLC3A2/4F2hc to form a functional heterodimeric complex that translocates small neutral L- and D-amino acids across the plasma membrane. Preferentially mediates exchange transport, but can also operate via facilitated diffusion (By similarity) (PubMed:<a href="http://www.uniprot.org/citations/10863037" target="\_blank">10863037</a>). Acts as a major transporter for glycine, L- and D-serine in the central nervous system. At the spinal cord and brainstem regulates glycine metabolism and glycinergic inhibitory neurotransmission by providing for glycine de novo synthesis from L- serine and glycine recycling from astrocytes to glycinergic motor neurons (By similarity). At Schaffer collateral-CA1 synapses mediates D-serine and glycine release that modulates post-synaptic activation of NMDA receptors and excitatory glutamatergic transmission (By similarity). May regulate D-serine release from mesenchymal progenitors located in developing subcutaneous adipose tissue, favoring white adipocyte over thermogenic beige adipocyte lineage commitment (By similarity).

### **Cellular Location**

Cell membrane {ECO:0000250|UniProtKB:P63115}; Multi-pass membrane protein.





Note=Colocalizes with OLIG2 in astrocytic processes. Localizes to the plasma membrane in mature adipocytes and to intracellular structures in preadipocytes {ECO:0000250|UniProtKB:P63115}

### **Tissue Location**

Expressed in brain, heart, kidney, liver, lung, pancreas, placenta, and skeletal muscle.

# AAA1 Antibody (C-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides

AAA1 Antibody (C-term) Blocking peptide - Images