

## CARD9 Antibody (N-term) Blocking peptide

Synthetic peptide Catalog # BP5985a

### **Specification**

## CARD9 Antibody (N-term) Blocking peptide - Product Information

Primary Accession Q9H257

Other Accession NP 434700.2, NP 434701.1

# CARD9 Antibody (N-term) Blocking peptide - Additional Information

**Gene ID** 64170

#### **Other Names**

Caspase recruitment domain-containing protein 9, hCARD9, CARD9

### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

### **Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

### CARD9 Antibody (N-term) Blocking peptide - Protein Information

Name CARD9 {ECO:0000303|PubMed:11053425, ECO:0000312|HGNC:HGNC:16391}

#### **Function**

Adapter protein that plays a key role in innate immune response against fungi by forming signaling complexes downstream of C- type lectin receptors (PubMed:<a href="http://www.uniprot.org/citations/26961233" target=" blank">26961233</a>, PubMed:<a href="http://www.uniprot.org/citations/33558980" target="blank">33558980</a>). CARD9mediated signals are essential for antifungal immunity against a subset of fungi from the phylum Ascomycota (PubMed: <a href="http://www.uniprot.org/citations/24231284" target="\_blank">24231284</a>, PubMed:<a href="http://www.uniprot.org/citations/25702837" target="blank">25702837</a>, PubMed:<a href="http://www.uniprot.org/citations/25057046" target="blank">25057046</a>, PubMed:<a href="http://www.uniprot.org/citations/26679537" target="\_blank">26679537</a>, PubMed:<a href="http://www.uniprot.org/citations/26961233" target="blank">26961233</a>, PubMed:<a href="http://www.uniprot.org/citations/26521038" target="blank">26521038</a>, PubMed:<a href="http://www.uniprot.org/citations/27777981" target="blank">27777981</a>, PubMed:<a href="http://www.uniprot.org/citations/29080677" target="blank">29080677</a>, PubMed:<a href="http://www.uniprot.org/citations/33558980" target="\_blank">33558980</a>). Transduces signals in myeloid cells downstream of C-type lectin receptors CLEC7A (dectin-1), CLEC6A (dectin-2) and CLEC4E (Mincle), which detect pathogenassociated molecular pattern metabolites (PAMPs), such as fungal carbohydrates, and trigger



CARD9 activation (By similarity). Upon activation, CARD9 homooligomerizes to form a nucleating helical template that recruits BCL10 via CARD-CARD interaction, thereby promoting polymerization of BCL10 and subsequent recruitment of MALT1: this leads to activation of NF-kappa-B and MAP kinase p38 (MAPK11, MAPK12, MAPK13 and/or MAPK14) pathways which stimulate expression of genes encoding pro-inflammatory cytokines and chemokines (PubMed:<a href="http://www.uniprot.org/citations/11053425" target="blank">11053425</a>, PubMed:<a

href="http://www.uniprot.org/citations/26488816" target="\_blank">26488816</a>, PubMed:<a href="http://www.uniprot.org/citations/31296852" target="\_blank">31296852</a>, PubMed:<a href="http://www.uniprot.org/citations/26961233" target="\_blank">26961233</a>, PubMed:<a href="http://www.uniprot.org/citations/26961233" target="\_blank">26961233</a>, PubMed:<a href="http://www.uniprot.org/citations/33558980" target="\_blank">33558980</a>). CARD9 signaling in antigen-presenting cells links innate sensing of fungi to the activation of adaptive immunity and provides a cytokine milieu that induces the development and subsequent of interleukin 17-producing T helper (Th17) cells (PubMed:<a

href="http://www.uniprot.org/citations/24231284" target="\_blank">24231284</a>). Also involved in activation of myeloid cells via classical ITAM- associated receptors and TLR: required for TLR-mediated activation of MAPK, while it is not required for TLR-induced activation of NF-kappa-B (By similarity). CARD9 can also be engaged independently of BCL10: forms a complex with RASGRF1 downstream of C-type lectin receptors, which recruits and activates HRAS, leading to ERK activation and the production of cytokines (By similarity). Acts as an important regulator of the intestinal commensal fungi (mycobiota) component of the gut microbiota (PubMed:<a href="http://www.uniprot.org/citations/33548172" target="\_blank">33548172</a>). Plays an essential role in antifungal immunity against dissemination of gut fungi: acts by promoting induction of antifungal IgG antibodies response in CX3CR1(+) macrophages to confer protection against disseminated C.albicans or C.auris infection (PubMed:<a

href="http://www.uniprot.org/citations/33548172" target="\_blank">33548172</a>). Also mediates immunity against other pathogens, such as certain bacteria, viruses and parasites; CARD9 signaling is however redundant with other innate immune responses (By similarity). In response to L.monocytogenes infection, required for the production of inflammatory cytokines activated by intracellular peptidoglycan: acts by connecting NOD2 recognition of peptidoglycan to downstream activation of MAP kinases (MAPK) without activating NF- kappa-B (By similarity).

Cellular Location Cytoplasm

#### **Tissue Location**

Expression is restricted to several populations of phagocytes, such as macrophages, monocytes, and dendritic cells (PubMed:33548172). Highly expressed in spleen (PubMed:11053425). Also detected in liver, placenta, lung, peripheral blood leukocytes and in brain (PubMed:11053425).

## CARD9 Antibody (N-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides

CARD9 Antibody (N-term) Blocking peptide - Images