

MAGEA10 Antibody (C-term) Blocking Peptide
Synthetic peptide
Catalog # BP6160a**Specification**

MAGEA10 Antibody (C-term) Blocking Peptide - Product Information

Primary Accession [P43363](#)
Other Accession [NP_066386](#)

MAGEA10 Antibody (C-term) Blocking Peptide - Additional Information

Gene ID 4109

Other Names

Melanoma-associated antigen 10, Cancer/testis antigen 110, CT110, MAGE-10 antigen, MAGEA10, MAGE10

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP6160a](/product/products/AP6160a) was selected from the C-term region of human MAGEA10. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

MAGEA10 Antibody (C-term) Blocking Peptide - Protein Information

Name MAGEA10

Synonyms MAGE10

Function

Not known, though may play a role in embryonal development and tumor transformation or aspects of tumor progression.

Cellular Location

Nucleus.

Tissue Location

Expressed in many tumors of several types, such as melanoma, head and neck squamous cell

carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for spermatogonia, spermatocytes and placenta.

MAGEA10 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

MAGEA10 Antibody (C-term) Blocking Peptide - Images

MAGEA10 Antibody (C-term) Blocking Peptide - Background

MAGEA10 is a member of the MAGEA gene family. The members of this family have their entire coding sequences located in the last exon, and the encoded proteins show 50 to 80% sequence identity between each other. The promoters and first exons of the MAGEA genes show considerable variability, suggesting that the existence of this gene family enables the same function to be expressed under different transcriptional controls. The MAGEA genes are expressed at a high level in a number of tumors of various histologic types, and are silent in normal tissues with the exception of testis and placenta. The MAGEA genes are clustered on chromosome Xq28. They may be implicated in some hereditary disorders, such as dyskeratosis congenita.

MAGEA10 Antibody (C-term) Blocking Peptide - References

Rogner, U.C., et al., Genomics 29(3):725-731 (1995). De Plaen, E., et al., Immunogenetics 40(5):360-369 (1994).