

**p53 Antibody (T55) Blocking peptide**  
**Synthetic peptide**  
**Catalog # BP6266d****Specification**

---

**p53 Antibody (T55) Blocking peptide - Product Information**Primary Accession [P04637](#)**p53 Antibody (T55) Blocking peptide - Additional Information****Gene ID** 7157**Other Names**

Cellular tumor antigen p53, Antigen NY-CO-13, Phosphoprotein p53, Tumor suppressor p53, TP53, P53

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP6266d](/products/AP6266d) was selected from the T55 region of human p53 Antibody (T55). A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**p53 Antibody (T55) Blocking peptide - Protein Information****Name** TP53**Synonyms** P53**Function**

Acts as a tumor suppressor in many tumor types; induces growth arrest or apoptosis depending on the physiological circumstances and cell type (PubMed: [11025664](http://www.uniprot.org/citations/11025664), PubMed: [12524540](http://www.uniprot.org/citations/12524540), PubMed: [12810724](http://www.uniprot.org/citations/12810724), PubMed: [15186775](http://www.uniprot.org/citations/15186775), PubMed: [15340061](http://www.uniprot.org/citations/15340061), PubMed: [17317671](http://www.uniprot.org/citations/17317671), PubMed: [17349958](http://www.uniprot.org/citations/17349958)).

[19556538](http://www.uniprot.org/citations/19556538), PubMed: [20673990](http://www.uniprot.org/citations/20673990), PubMed: [20959462](http://www.uniprot.org/citations/20959462), PubMed: [22726440](http://www.uniprot.org/citations/22726440), PubMed: [24051492](http://www.uniprot.org/citations/24051492), PubMed: [9840937](http://www.uniprot.org/citations/9840937), PubMed: [24652652](http://www.uniprot.org/citations/24652652)). Involved in cell cycle regulation as a trans-activator that acts to negatively regulate cell division by controlling a set of genes required for this process (PubMed: [11025664](http://www.uniprot.org/citations/11025664), PubMed: [12524540](http://www.uniprot.org/citations/12524540), PubMed: [12810724](http://www.uniprot.org/citations/12810724), PubMed: [15186775](http://www.uniprot.org/citations/15186775), PubMed: [15340061](http://www.uniprot.org/citations/15340061), PubMed: [17317671](http://www.uniprot.org/citations/17317671), PubMed: [17349958](http://www.uniprot.org/citations/17349958), PubMed: [19556538](http://www.uniprot.org/citations/19556538), PubMed: [20673990](http://www.uniprot.org/citations/20673990), PubMed: [20959462](http://www.uniprot.org/citations/20959462), PubMed: [22726440](http://www.uniprot.org/citations/22726440), PubMed: [24051492](http://www.uniprot.org/citations/24051492), PubMed: [9840937](http://www.uniprot.org/citations/9840937), PubMed: [24652652](http://www.uniprot.org/citations/24652652)). One of the activated genes is an inhibitor of cyclin-dependent kinases. Apoptosis induction seems to be mediated either by stimulation of BAX and FAS antigen expression, or by repression of Bcl-2 expression. Its pro-apoptotic activity is activated via its interaction with PPP1R13B/ASPP1 or TP53BP2/ASPP2 (PubMed: [12524540](http://www.uniprot.org/citations/12524540)). However, this activity is inhibited when the interaction with PPP1R13B/ASPP1 or TP53BP2/ASPP2 is displaced by PPP1R13L/iASPP (PubMed: [12524540](http://www.uniprot.org/citations/12524540)). In cooperation with mitochondrial PPIF is involved in activating oxidative stress-induced necrosis; the function is largely independent of transcription. Induces the transcription of long intergenic non-coding RNA p21 (lincRNA-p21) and lincRNA-Mkln1. LincRNA-p21 participates in TP53-dependent transcriptional repression leading to apoptosis and seems to have an effect on cell-cycle regulation. Implicated in Notch signaling cross-over. Prevents CDK7 kinase activity when associated to CAK complex in response to DNA damage, thus stopping cell cycle progression. Isoform 2 enhances the transactivation activity of isoform 1 from some but not all TP53-inducible promoters. Isoform 4 suppresses transactivation activity and impairs growth suppression mediated by isoform 1. Isoform 7 inhibits isoform 1-mediated apoptosis. Regulates the circadian clock by repressing CLOCK-BMAL1-mediated transcriptional activation of PER2 (PubMed: [24051492](http://www.uniprot.org/citations/24051492)).

### Cellular Location

Cytoplasm. Nucleus. Nucleus, PML body Endoplasmic reticulum. Mitochondrion matrix. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome Note=Recruited into PML bodies together with CHEK2 (PubMed:12810724) Translocates to mitochondria upon oxidative stress (PubMed:22726440) Translocates to mitochondria in response to mitomycin C treatment (PubMed:27323408). Competitive inhibition of TP53 interaction with HSPA9/MOT-2 by UBXN2A results in increased protein abundance and subsequent translocation of TP53 to the nucleus (PubMed:24625977) [Isoform 2]: Nucleus. Cytoplasm. Note=Localized mainly in the nucleus with minor staining in the cytoplasm [Isoform 4]: Nucleus. Cytoplasm. Note=Predominantly nuclear but translocates to the cytoplasm following cell stress [Isoform 8]: Nucleus. Cytoplasm. Note=Localized in both nucleus and cytoplasm in most cells. In some cells, forms foci in the nucleus that are different from nucleoli

### Tissue Location

Ubiquitous. Isoforms are expressed in a wide range of normal tissues but in a tissue-dependent

manner. Isoform 2 is expressed in most normal tissues but is not detected in brain, lung, prostate, muscle, fetal brain, spinal cord and fetal liver. Isoform 3 is expressed in most normal tissues but is not detected in lung, spleen, testis, fetal brain, spinal cord and fetal liver. Isoform 7 is expressed in most normal tissues but is not detected in prostate, uterus, skeletal muscle and breast. Isoform 8 is detected only in colon, bone marrow, testis, fetal brain and intestine. Isoform 9 is expressed in most normal tissues but is not detected in brain, heart, lung, fetal liver, salivary gland, breast or intestine

### **p53 Antibody (T55) Blocking peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

### **p53 Antibody (T55) Blocking peptide - Images**

### **p53 Antibody (T55) Blocking peptide - Background**

Tumor protein p53, a nuclear protein, plays an essential role in the regulation of cell cycle, specifically in the transition from G0 to G1. It is found in very low levels in normal cells, however, in a variety of transformed cell lines, it is expressed in high amounts, and believed to contribute to transformation and malignancy. p53 is a DNA-binding protein containing DNA-binding, oligomerization and transcription activation domains. It is postulated to bind as a tetramer to a p53-binding site and activate expression of downstream genes that inhibit growth and/or invasion, and thus function as a tumor suppressor. Mutants of p53 that frequently occur in a number of different human cancers fail to bind the consensus DNA binding site, and hence cause the loss of tumor suppressor activity. Alterations of the TP53 gene occur not only as somatic mutations in human malignancies, but also as germline mutations in some cancer-prone families with Li-Fraumeni syndrome.

### **p53 Antibody (T55) Blocking peptide - References**

Blanchette, P., et al., Mol. Cell. Biol. 24(21):9619-9629 (2004). Adachi, K., et al., Oncogene 23(47):7791-7798 (2004). Zhang, Y., et al., J. Biol. Chem. 279(41):42545-42551 (2004). Anazawa, Y., et al., Oncogene 23(46):7621-7627 (2004). Montagnoli, A., et al., Cancer Res. 64(19):7110-7116 (2004).