

AGR2 Antibody (Center) Blocking Peptide
Synthetic peptide
Catalog # BP6279c**Specification**

AGR2 Antibody (Center) Blocking Peptide - Product InformationPrimary Accession [O95994](#)**AGR2 Antibody (Center) Blocking Peptide - Additional Information****Gene ID** 10551**Other Names**

Anterior gradient protein 2 homolog, AG-2, hAG-2, HPC8, Secreted cement gland protein XAG-2 homolog, AGR2, AG2

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP6279c](/product/products/AP6279c) was selected from the Center region of human AGR2. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

AGR2 Antibody (Center) Blocking Peptide - Protein Information**Name** AGR2**Synonyms** AG2**Function**

Required for MUC2 post-transcriptional synthesis and secretion. May play a role in the production of mucus by intestinal cells (By similarity). Proto-oncogene that may play a role in cell migration, cell differentiation and cell growth. Promotes cell adhesion (PubMed:[23274113](http://www.uniprot.org/citations/23274113)).

Cellular Location

Secreted. Endoplasmic reticulum {ECO:0000250|UniProtKB:O88312}

Tissue Location

Expressed strongly in trachea, lung, stomach, colon, prostate and small intestine. Expressed weakly in pituitary gland, salivary gland, mammary gland, bladder, appendix, ovary, fetal lung, uterus, pancreas, kidney, fetal kidney, testis, placenta, thyroid gland and in estrogen receptor (ER)-positive breast cancer cell lines

AGR2 Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

AGR2 Antibody (Center) Blocking Peptide - Images

AGR2 Antibody (Center) Blocking Peptide - Background

Anterior gradient 2 (AGR2) is known as a cancer cell marker specifically up-regulated in response to depletion of serum and oxygen. AGR2 has been identified as a tumor marker in primary and secondary cancer lesions, and as a marker for detection of circulating tumor cells (CTCs). Elevated levels of AGR2 are known to increase the metastatic potential of cancer cells, but conditions leading to increased expression of AGR2 are not well understood.

AGR2 Antibody (Center) Blocking Peptide - References

Zweitzig,D.R., Mol. Cell. Biochem. 306 (1-2), 255-260 (2007)Zhang,Y., Prostate Cancer Prostatic Dis. 10 (3), 293-300 (2007)Fletcher,G.C., Br. J. Cancer 88 (4), 579-585 (2003)