

# PARP16 Antibody (C-term) Blocking Peptide

Synthetic peptide Catalog # BP6299d

## **Specification**

# PARP16 Antibody (C-term) Blocking Peptide - Product Information

Primary Accession

# PARP16 Antibody (C-term) Blocking Peptide - Additional Information

**Gene ID 54956** 

#### **Other Names**

Mono [ADP-ribose] polymerase PARP16, ADP-ribosyltransferase diphtheria toxin-like 15, Poly [ADP-ribose] polymerase 16, PARP-16, PAR16

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Q8N5Y8** 

## **Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

## PARP16 Antibody (C-term) Blocking Peptide - Protein Information

Name PARP16 {ECO:0000303|PubMed:20106667, ECO:0000312|HGNC:HGNC:26040}

# **Function**

Intracellular mono-ADP-ribosyltransferase that plays a role in different processes, such as protein translation and unfolded protein response (UPR), through the mono-ADP-ribosylation of proteins involved in those processes (PubMed: <a href="http://www.uniprot.org/citations/22701565" target=" blank">22701565</a>, PubMed:<a href="http://www.uniprot.org/citations/23103912" target="blank">23103912</a>, PubMed:<a href="http://www.uniprot.org/citations/25043379" target="blank">25043379</a>, PubMed:<a href="http://www.uniprot.org/citations/34314702" target="blank">34314702</a>). Acts as an inhibitor of protein translation by catalyzing mono-ADP-ribosylation of ribosomal subunits, such as RPL14 and RPS6, thereby inhibiting polysome assembly and mRNA loading (PubMed: <a href="http://www.uniprot.org/citations/34314702" target=" blank">34314702</a>). Mono-ADP-ribosylation of ribosomal subunits is promoted by NMNAT2 (PubMed: <a href="http://www.uniprot.org/citations/34314702" target=" blank">34314702</a>). Involved in the unfolded protein response (UPR) by ADP-ribosylating and activating EIF2AK3 and ERN1, two important UPR effectors (PubMed: <a href="http://www.uniprot.org/citations/23103912" target=" blank">23103912</a>). May also mediate mono-ADP-ribosylation of karyopherin KPNB1 a nuclear import factor (PubMed:<a href="http://www.uniprot.org/citations/22701565" target=" blank">22701565</a>). May not modify proteins on arginine or cysteine residues



compared to other mono-ADP-ribosyltransferases (PubMed:<a href="http://www.uniprot.org/citations/22701565" target=" blank">22701565</a>).

#### **Cellular Location**

Endoplasmic reticulum membrane; Single-pass type IV membrane protein

# PARP16 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

## • Blocking Peptides

PARP16 Antibody (C-term) Blocking Peptide - Images

PARP16 Antibody (C-term) Blocking Peptide - Background

The function of this protein has not been specifically defined.

# PARP16 Antibody (C-term) Blocking Peptide - References

Ame, J.C., et al. Bioessays 26(8):882-893(2004)