

# Kinesin 5A (Kif5A) Antibody (C-term) Blocking peptide

Synthetic peptide Catalog # BP6349b

# **Specification**

# Kinesin 5A (Kif5A) Antibody (C-term) Blocking peptide - Product Information

**Primary Accession** 

**Q12840** 

# Kinesin 5A (Kif5A) Antibody (C-term) Blocking peptide - Additional Information

**Gene ID 3798** 

#### **Other Names**

Kinesin heavy chain isoform 5A, Kinesin heavy chain neuron-specific 1, Neuronal kinesin heavy chain, NKHC, KIF5A, NKHC1

# **Target/Specificity**

The synthetic peptide sequence used to generate the antibody <a href=/product/products/AP6349b>AP6349b</a> was selected from the C-term region of human Kif5A. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

### Kinesin 5A (Kif5A) Antibody (C-term) Blocking peptide - Protein Information

Name KIF5A

Synonyms NKHC1

### **Function**

Microtubule-dependent motor required for slow axonal transport of neurofilament proteins (NFH, NFM and NFL). Can induce formation of neurite-like membrane protrusions in non-neuronal cells in a ZFYVE27-dependent manner. The ZFYVE27-KIF5A complex contributes to the vesicular transport of VAPA, VAPB, SURF4, RAB11A, RAB11B and RTN3 proteins in neurons. Required for anterograde axonal transportation of MAPK8IP3/JIP3 which is essential for MAPK8IP3/JIP3 function in axon elongation.

#### **Cellular Location**

Cytoplasm, perinuclear region {ECO:0000250|UniProtKB:Q6QLM7}. Cytoplasm, cytoskeleton



Tel: 858.875.1900 Fax: 858.875.1999

{ECO:0000250|UniProtKB:Q6QLM7}. Perikaryon {ECO:0000250|UniProtKB:Q6QLM7}. Note=Concentrated in the cell body of the neurons, particularly in the perinuclear region {ECO:0000250|UniProtKB:Q6QLM7}

#### **Tissue Location**

Distributed throughout the CNS but is highly enriched in subsets of neurons

# Kinesin 5A (Kif5A) Antibody (C-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

# Blocking Peptides

Kinesin 5A (Kif5A) Antibody (C-term) Blocking peptide - Images

### Kinesin 5A (Kif5A) Antibody (C-term) Blocking peptide - Background

Kif5A is a member of the kinesin family of proteins. Members of this family are part of a multisubunit complex that functions as a microtubule motor in intracellular organelle transport. Mutations in this protein cause autosomal dominant spastic paraplegia 10.

# Kinesin 5A (Kif5A) Antibody (C-term) Blocking peptide - References

Fichera, M., et al., Neurology 63(6):1108-1110 (2004). Reid, E., et al., Am. J. Hum. Genet. 71(5):1189-1194 (2002).Kanai, Y., et al., J. Neurosci. 20(17):6374-6384 (2000).Reid, E., et al., Am. J. Hum. Genet. 65(3):757-763 (1999). Rahman, A., et al., J. Cell Biol. 146(6):1277-1288 (1999).