

**PLD1 Antibody (N-term) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP6377a****Specification**

---

**PLD1 Antibody (N-term) Blocking Peptide - Product Information**Primary Accession [Q13393](#)**PLD1 Antibody (N-term) Blocking Peptide - Additional Information**

Gene ID 5337

**Other Names**

Phospholipase D1, PLD 1, hPLD1, Choline phosphatase 1, Phosphatidylcholine-hydrolyzing phospholipase D1, PLD1

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP6377a](/products/AP6377a) was selected from the N-term region of human PLD1. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**PLD1 Antibody (N-term) Blocking Peptide - Protein Information**Name PLD1 ([HGNC:9067](#))**Function**

Function as phospholipase selective for phosphatidylcholine (PubMed: [8530346](http://www.uniprot.org/citations/8530346), PubMed: [9582313](http://www.uniprot.org/citations/9582313), PubMed: [25936805](http://www.uniprot.org/citations/25936805)). Implicated as a critical step in numerous cellular pathways, including signal transduction, membrane trafficking, and the regulation of mitosis. May be involved in the regulation of perinuclear intravesicular membrane traffic (By similarity).

**Cellular Location**

Cytoplasm, perinuclear region {ECO:0000250|UniProtKB:Q9Z280}. Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:Q9Z280}; Lipid-anchor {ECO:0000250|UniProtKB:Q9Z280};

Cytoplasmic side {ECO:0000250|UniProtKB:Q9Z280}. Golgi apparatus membrane {ECO:0000250|UniProtKB:Q9Z280}; Lipid-anchor {ECO:0000250|UniProtKB:Q9Z280}; Cytoplasmic side {ECO:0000250|UniProtKB:Q9Z280}. Late endosome membrane {ECO:0000250|UniProtKB:Q9Z280}; Lipid-anchor {ECO:0000250|UniProtKB:Q9Z280}; Cytoplasmic side {ECO:0000250|UniProtKB:Q9Z280}

**Tissue Location**

Expressed abundantly in the pancreas and heart and at high levels in brain, placenta, spleen, uterus and small intestine

**PLD1 Antibody (N-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**PLD1 Antibody (N-term) Blocking Peptide - Images****PLD1 Antibody (N-term) Blocking Peptide - Background**

Phosphatidylcholine (PC)-specific phospholipases D (PLDs; EC 3.1.4.4) catalyze the hydrolysis of PC to produce phosphatidic acid and choline. A range of agonists acting through G protein-coupled receptors and receptor tyrosine kinases stimulate this hydrolysis. PC-specific PLD activity has been implicated in numerous cellular pathways, including signal transduction, membrane trafficking, and the regulation of mitosis (Hammond et al., 1995 [PubMed 8530346]).[supplied by OMIM]

**PLD1 Antibody (N-term) Blocking Peptide - References**

Sun,Y., Proc. Natl. Acad. Sci. U.S.A. 105 (24), 8286-8291 (2008)Sethu,S., J. Immunol. 180 (9), 6027-6034 (2008)Nagasaki,A., Cell Struct. Funct. 33 (1), 27-33 (2008)