

Synphilin-1 (SNCAIP) Antibody (C-term) Blocking Peptide
Synthetic peptide
Catalog # BP6411b**Specification**

Synphilin-1 (SNCAIP) Antibody (C-term) Blocking Peptide - Product InformationPrimary Accession
Other Accession[O9Y6H5](#)
[NP_005451](#)**Synphilin-1 (SNCAIP) Antibody (C-term) Blocking Peptide - Additional Information****Gene ID** 9627**Other Names**

Synphilin-1, Sph1, Alpha-synuclein-interacting protein, SNCAIP

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP6411b](/product/products/AP6411b) was selected from the SNCAIP region of human Synphilin-1 (SNCAIP). A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

Synphilin-1 (SNCAIP) Antibody (C-term) Blocking Peptide - Protein Information**Name** SNCAIP**Function**

Isoform 2 inhibits the ubiquitin ligase activity of SIAH1 and inhibits proteasomal degradation of target proteins. Isoform 2 inhibits autoubiquitination and proteasomal degradation of SIAH1, and thereby increases cellular levels of SIAH. Isoform 2 modulates SNCA monoubiquitination by SIAH1.

Cellular Location

Cytoplasm. Note=Detected in cytoplasmic inclusion bodies, together with SNCA

Tissue Location

Detected in brain (at protein level). Widely expressed, with highest levels in brain, heart and placenta

Synphilin-1 (SNCAIP) Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

Synphilin-1 (SNCAIP) Antibody (C-term) Blocking Peptide - Images

Synphilin-1 (SNCAIP) Antibody (C-term) Blocking Peptide - Background

Parkinson is the second most common neurodegenerative disease after Alzheimers. About 1 percent of people over the age of 65 and 3 percent of people over the age of 75 are affected by the disease. The mutation is the most common cause of Parkinson disease identified to date. Synuclein alpha interacting protein (Synphilin-1) contains several protein-protein interaction domains and interacts with alpha synuclein in neurons. Mutations of SNCAIP have been linked to Parkinson disease. The amino acid sequence of synphilin-1 shows extensive homology with its human counterpart, especially in regions containing ankyrin-like motifs and the coiled-coil domain. Expression of mouse synphilin-1 in tissues is similar to its human counterpart. Synphilin-1 has an important role in the formation of aggregates and cytotoxicity in Parkinson disease and that Dornfin may be involved in the pathogenic process by ubiquitylation of synphilin-1. Role of synphilin-1 in synaptic function and protein degradation and in the molecular mechanisms leading to neurodegeneration in Parkinson disease

Synphilin-1 (SNCAIP) Antibody (C-term) Blocking Peptide - References

Kruger,R. Cell Tissue Res. 318 (1), 195-199 (2004)Lee,G., etal. J. Biol. Chem. 279 (8), 6834-6839 (2004)Tanaka,M., et al. J. Biol. Chem. 279 (6), 4625-4631 (2004)Nagano,Y., et al. J. Biol. Chem. 278 (51), 51504-51514 (2003)Marx,F.P., etal. Hum. Mol. Genet. 12 (11), 1223-1231 (2003)Junn,E., et al. J. Biol. Chem. 277 (49), 47870-47877 (2002)Chung,K.K., et al. Nat. Med. 7 (10), 1144-1150 (2001)Kawamata,H., et al. J. Neurochem. 77 (3), 929-934 (2001)Engelender,S., et al. Nat. Genet. 22 (1), 110-114 (1999)