

# COPEB Antibody (N-term) Blocking Peptide

Synthetic peptide Catalog # BP6588a

# Specification

# **COPEB Antibody (N-term) Blocking Peptide - Product Information**

Primary Accession

<u>Q99612</u>

# **COPEB** Antibody (N-term) Blocking Peptide - Additional Information

Gene ID 1316

#### **Other Names**

Krueppel-like factor 6, B-cell-derived protein 1, Core promoter element-binding protein, GC-rich sites-binding factor GBF, Proto-oncogene BCD1, Suppressor of tumorigenicity 12 protein, Transcription factor Zf9, KLF6, BCD1, COPEB, CPBP, ST12

## Target/Specificity

The synthetic peptide sequence used to generate the antibody <a href=/products/AP6588a>AP6588a</a> was selected from the N-term region of human COPEB. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions** 

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

## **COPEB Antibody (N-term) Blocking Peptide - Protein Information**

Name KLF6

Synonyms BCD1, COPEB, CPBP, ST12

Function

Transcriptional activator (By similarity). Binds a GC box motif. Could play a role in B-cell growth and development.

**Cellular Location** Nucleus.

#### **Tissue Location**

Highly expressed in placenta followed by spleen, thymus, prostate, testis, small intestine and



colon. Weakly expressed in pancreas, lung, liver, heart and skeletal muscle. Also expressed in fetal brain, spleen and thymus

# **COPEB Antibody (N-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

<u>Blocking Peptides</u>

## **COPEB Antibody (N-term) Blocking Peptide - Images**

#### **COPEB Antibody (N-term) Blocking Peptide - Background**

COPEB is a member of the Kruppel-like family of transcription factors. The zinc finger protein is a transcriptional activator, and functions as a tumor suppressor.

#### **COPEB Antibody (N-term) Blocking Peptide - References**

Sangodkar, J., Eur. J. Cancer 45 (4), 666-676 (2009) DiFeo, A., Drug Resist. Updat. 12 (1-2), 1-7 (2009)