

TORC1 Antibody (N-term) Blocking Peptide
Synthetic peptide
Catalog # BP6968a**Specification**

TORC1 Antibody (N-term) Blocking Peptide - Product InformationPrimary Accession [Q6UUV9](#)**TORC1 Antibody (N-term) Blocking Peptide - Additional Information****Gene ID** 23373**Other Names**

CREB-regulated transcription coactivator 1, Mucoepidermoid carcinoma translocated protein 1, Transducer of regulated cAMP response element-binding protein 1, TORC-1, Transducer of CREB protein 1, CRTC1 (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=16062)
HGNC:16062

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP6968a](/products/AP6968a) was selected from the N-term region of human TORC1. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

TORC1 Antibody (N-term) Blocking Peptide - Protein Information**Name** CRTC1 ([HGNC:16062](#))**Function**

Transcriptional coactivator for CREB1 which activates transcription through both consensus and variant cAMP response element (CRE) sites. Acts as a coactivator, in the SIK/TORC signaling pathway, being active when dephosphorylated and acts independently of CREB1 'Ser-133' phosphorylation. Enhances the interaction of CREB1 with TAF4. Regulates the expression of specific CREB-activated genes such as the steroidogenic gene, StAR. Potent coactivator of PGC1alpha and inducer of mitochondrial biogenesis in muscle cells. In the hippocampus, involved in late-phase long-term potentiation (L-LTP) maintenance at the Schaffer collateral-CA1 synapses. May be required for dendritic growth of developing cortical neurons (By similarity). In concert with

SIK1, regulates the light-induced entrainment of the circadian clock. In response to light stimulus, coactivates the CREB-mediated transcription of PER1 which plays an important role in the photic entrainment of the circadian clock.

Cellular Location

Cytoplasm. Nucleus. Note=Cytoplasmic when phosphorylated by SIK or AMPK and when sequestered by 14-3-3 proteins (PubMed:16817901) Translocated to the nucleus on Ser-151 dephosphorylation, instigated by a number of factors including calcium ion and cAMP levels (PubMed:15589160). Light stimulation triggers a nuclear accumulation in the suprachiasmatic nucleus (SCN) of the brain (By similarity) {ECO:0000250|UniProtKB:Q68ED7, ECO:0000269|PubMed:15589160, ECO:0000269|PubMed:16817901}

Tissue Location

Highly expressed in adult and fetal brain. Located to specific regions such as the prefrontal cortex and cerebellum. Very low expression in other tissues such as heart, spleen, lung, skeletal muscle, salivary gland, ovary and kidney.

TORC1 Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

TORC1 Antibody (N-term) Blocking Peptide - Images**TORC1 Antibody (N-term) Blocking Peptide - Background**

TORC1 is the transcriptional coactivator for CREB1 which activates transcription through both consensus and variant cAMP response element (CRE) sites. It Acts as a coactivator, in the SIK/TORC signaling pathway, being active when dephosphorylated and acts independently of CREB1 'Ser-133' phosphorylation and Enhances the interaction of CREB1 with TAF4. It regulates the expression of specific CREB-activated genes such as the steroidogenic gene, StAR.

TORC1 Antibody (N-term) Blocking Peptide - References

Grimwood,J., et.al., Nature 428 (6982), 529-535 (2004)