

PDK4 Antibody (C-term) Blocking Peptide
Synthetic peptide
Catalog # BP7041b**Specification**

PDK4 Antibody (C-term) Blocking Peptide - Product InformationPrimary Accession [Q16654](#)**PDK4 Antibody (C-term) Blocking Peptide - Additional Information****Gene ID** 5166**Other Names**

[Pyruvate dehydrogenase (acetyl-transferring)] kinase isozyme 4, mitochondrial, Pyruvate dehydrogenase kinase isoform 4, PDK4, PDHK4

Target/Specificity

The synthetic peptide sequence is selected from aa 396-410 of human PDK4.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

PDK4 Antibody (C-term) Blocking Peptide - Protein Information**Name** PDK4**Synonyms** PDHK4**Function**

Kinase that plays a key role in regulation of glucose and fatty acid metabolism and homeostasis via phosphorylation of the pyruvate dehydrogenase subunits PDHA1 and PDHA2. This inhibits pyruvate dehydrogenase activity, and thereby regulates metabolite flux through the tricarboxylic acid cycle, down-regulates aerobic respiration and inhibits the formation of acetyl-coenzyme A from pyruvate. Inhibition of pyruvate dehydrogenase decreases glucose utilization and increases fat metabolism in response to prolonged fasting and starvation. Plays an important role in maintaining normal blood glucose levels under starvation, and is involved in the insulin signaling cascade. Via its regulation of pyruvate dehydrogenase activity, plays an important role in maintaining normal blood pH and in preventing the accumulation of ketone bodies under starvation. In the fed state, mediates cellular responses to glucose levels and to a high-fat diet. Regulates both fatty acid oxidation and de novo fatty acid biosynthesis. Plays a role in the generation of reactive oxygen species. Protects detached epithelial cells against anoikis. Plays a

role in cell proliferation via its role in regulating carbohydrate and fatty acid metabolism.

Cellular Location

Mitochondrion matrix.

Tissue Location

Ubiquitous; highest levels of expression in heart and skeletal muscle.

PDK4 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

PDK4 Antibody (C-term) Blocking Peptide - Images**PDK4 Antibody (C-term) Blocking Peptide - Background**

PDK4 inhibits the mitochondrial pyruvate dehydrogenase complex by phosphorylation of the E1 alpha subunit, thus contributing to the regulation of glucose metabolism.

PDK4 Antibody (C-term) Blocking Peptide - References

Blume-Jensen P, et al. Nature 2001. 411: 355. Cantrell D, J. Cell Sci. 2001. 114: 1439. Jhian S. Oncogene 2000. 19: 5590. Manning G, et al. Science 2002. 298: 1912. Moller, D, et al. Am. J. Physiol. 1994. 266: C351-C359. Robertson, S. et al. Trends Genet. 2000. 16: 368. Robinson D, et al. Oncogene 2000. 19: 5548. Van der Ven, P, et al. Hum. Molec. Genet. 1993. 2: 1889. Vanhaesebroeck, B, et al. Biochem. J. 2000. 346: 561. Van Weering D, et al. Recent Results Cancer Res. 1998. 154: 271.

PDK4 Antibody (C-term) Blocking Peptide - Citations

- [A mechanical trigger for the trot-gallop transition in horses.](#)