

**STK38 Antibody (C-term) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP7074b****Specification**

---

**STK38 Antibody (C-term) Blocking Peptide - Product Information**Primary Accession [Q15208](#)**STK38 Antibody (C-term) Blocking Peptide - Additional Information****Gene ID** 11329**Other Names**Serine/threonine-protein kinase 38, NDR1 protein kinase, Nuclear Dbf2-related kinase 1, STK38  
{ECO:0000312|EMBL:AAH120851}**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP7074b](/product/products/AP7074b) was selected from the C-term region of human STK38. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**STK38 Antibody (C-term) Blocking Peptide - Protein Information****Name** STK38 {ECO:0000312|EMBL:AAH12085.1}**Function**

Negative regulator of MAP3K1/2 signaling. Converts MAP3K2 from its phosphorylated form to its non-phosphorylated form and inhibits autophosphorylation of MAP3K2.

**Cellular Location**

Nucleus. Cytoplasm.

**Tissue Location**

Ubiquitously expressed with highest levels observed in peripheral blood leukocytes.

## **STK38 Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

## **STK38 Antibody (C-term) Blocking Peptide - Images**

## **STK38 Antibody (C-term) Blocking Peptide - Background**

Stk38, also known as NUCLEAR DBF2-RELATED PROTEIN, contains 1 protein kinase domain that interacts with mob1 and mob2. The homodimeric s100b binds to two molecules of stk38. It is ubiquitously expressed with highest levels observed in peripheral blood leukocytes. Stk38 is activated by binding of s100b which releases autoinhibitory n-lobe interactions, enabling ATP to bind and the autophosphorylation of ser-281. thr-444 then undergoes calcium-dependent phosphorylation by an upstream kinase. Interactions between phosphorylated thr-444 and the n-lobe promote additional structural changes that complete the activation of the kinase. Autoinhibition is also released by the binding of mob1/mobkl1a and mob2/hcca2 to the n-terminal of stk38.

## **STK38 Antibody (C-term) Blocking Peptide - References**

Devroe, E., et al., J. Biol. Chem. 279(23):24444-24451 (2004). Tamaskovic, R., et al., J. Biol. Chem. 278(9):6710-6718 (2003). Tripodis, N., et al., Genome Res. 8(6):631-643 (1998). Millward, T., et al., Proc. Natl. Acad. Sci. U.S.A. 92(11):5022-5026 (1995).