

ROCK1 Antibody (C-term) Blocking peptide
Synthetic peptide
Catalog # BP7096b**Specification**

ROCK1 Antibody (C-term) Blocking peptide - Product InformationPrimary Accession [Q13464](#)**ROCK1 Antibody (C-term) Blocking peptide - Additional Information****Gene ID** 6093**Other Names**

Rho-associated protein kinase 1, Renal carcinoma antigen NY-REN-35, Rho-associated, coiled-coil-containing protein kinase 1, Rho-associated, coiled-coil-containing protein kinase I, ROCK-I, p160 ROCK-1, p160ROCK, ROCK1

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP7096b](/product/products/AP7096b) was selected from the region of human ROCK1 C-term. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

ROCK1 Antibody (C-term) Blocking peptide - Protein Information**Name** ROCK1**Function**

Protein kinase which is a key regulator of the actin cytoskeleton and cell polarity (PubMed:[10436159](http://www.uniprot.org/citations/10436159), PubMed:[10652353](http://www.uniprot.org/citations/10652353), PubMed:[11018042](http://www.uniprot.org/citations/11018042), PubMed:[11283607](http://www.uniprot.org/citations/11283607), PubMed:[17158456](http://www.uniprot.org/citations/17158456), PubMed:[18573880](http://www.uniprot.org/citations/18573880), PubMed:[19131646](http://www.uniprot.org/citations/19131646), PubMed:[8617235](http://www.uniprot.org/citations/8617235), PubMed:[9722579](http://www.uniprot.org/citations/9722579)). Involved in

regulation of smooth muscle contraction, actin cytoskeleton organization, stress fiber and focal adhesion formation, neurite retraction, cell adhesion and motility via phosphorylation of DAPK3, GFAP, LIMK1, LIMK2, MYL9/MLC2, TPPP, PFN1 and PPP1R12A (PubMed:10436159, PubMed:10652353, PubMed:11018042, PubMed:11283607, PubMed:17158456, PubMed:18573880, PubMed:19131646, PubMed:8617235, PubMed:9722579, PubMed:23093407, PubMed:23355470). Phosphorylates FHOD1 and acts synergistically with it to promote SRC-dependent non-apoptotic plasma membrane blebbing (PubMed:18694941). Phosphorylates JIP3 and regulates the recruitment of JNK to JIP3 upon UVB-induced stress (PubMed:19036714). Acts as a suppressor of inflammatory cell migration by regulating PTEN phosphorylation and stability (By similarity). Acts as a negative regulator of VEGF-induced angiogenic endothelial cell activation (PubMed:19181962). Required for centrosome positioning and centrosome-dependent exit from mitosis (By similarity). Plays a role in terminal erythroid differentiation (PubMed:21072057). Inhibits podocyte motility via regulation of actin cytoskeletal dynamics and phosphorylation of CFL1 (By similarity). Promotes keratinocyte terminal differentiation (PubMed:19997641). Involved in osteoblast compaction through the fibronectin fibrillogenesis cell-mediated matrix assembly process, essential for osteoblast mineralization (By similarity). May regulate closure of the eyelids and ventral body wall by inducing the assembly of actomyosin bundles (By similarity).

Cellular Location

Cytoplasm. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole {ECO:0000250|UniProtKB:P70335}. Golgi apparatus membrane; Peripheral membrane protein. Cell projection, bleb. Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:P70335}. Cell membrane {ECO:0000250|UniProtKB:P70335}. Cell projection, lamellipodium {ECO:0000250|UniProtKB:P70335}. Cell projection, ruffle {ECO:0000250|UniProtKB:P70335}. Note=A small proportion is associated with Golgi membranes (PubMed:12773565). Associated with the mother centriole and an intercentriolar linker (By similarity). Colocalizes with ITGB1BP1 and ITGB1 at the cell membrane predominantly in lamellipodia and membrane ruffles, but also in retraction fibers (By similarity). Localizes at the cell membrane in an ITGB1BP1-dependent manner (By similarity). {ECO:0000250|UniProtKB:P70335, ECO:0000269|PubMed:12773565}

Tissue Location

Detected in blood platelets.

ROCK1 Antibody (C-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

ROCK1 Antibody (C-term) Blocking peptide - Images

ROCK1 Antibody (C-term) Blocking peptide - Background

ROCK1 is a protein serine/threonine kinase that is activated when bound to the GTP-bound form of Rho. The small GTPase Rho regulates formation of focal adhesions and stress fibers of fibroblasts, as well as adhesion and aggregation of platelets and lymphocytes by shuttling between the inactive GDP-bound form and the active GTP-bound form. Rho is also essential in cytokinesis and plays a role in transcriptional activation by serum response factor. ROCK, a downstream effector of Rho, phosphorylates and activates LIM kinase, which in turn, phosphorylates cofilin, inhibiting its actin-depolymerizing activity.

ROCK1 Antibody (C-term) Blocking peptide - References

Mammoto, A., et al., J. Biol. Chem. 279(25):26323-26330 (2004). Oi, K., et al., Arterioscler. Thromb. Vasc. Biol. 24(5):918-922 (2004). Wang, D.S., et al., World J. Gastroenterol. 10(2):299-302 (2004). Dvorsky, R., et al., J. Biol. Chem. 279(8):7098-7104 (2004). Wozniak, M.A., et al., J. Cell Biol. 163(3):583-595 (2003).