

Catalog # BP7114b

**BIKE Antibody (C-term) Blocking Peptide** Synthetic peptide

# Specification

# BIKE Antibody (C-term) Blocking Peptide - Product Information

Primary Accession

<u>Q9NSY1</u>

# BIKE Antibody (C-term) Blocking Peptide - Additional Information

Gene ID 55589

Other Names BMP-2-inducible protein kinase, BIKe, BMP2K, BIKE

Target/Specificity

The synthetic peptide sequence used to generate the antibody <a href=/product/products/AP7114b>AP7114b</a> was selected from the C-term region of human BIKE. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage** Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

### **Precautions** This product is for research use only. Not for use in diagnostic or therapeutic procedures.

### BIKE Antibody (C-term) Blocking Peptide - Protein Information

Name BMP2K

Synonyms BIKE

**Function** May be involved in osteoblast differentiation.

Cellular Location Nucleus.

### **BIKE Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.



#### <u>Blocking Peptides</u>

# BIKE Antibody (C-term) Blocking Peptide - Images

### BIKE Antibody (C-term) Blocking Peptide - Background

BIKE is the human homolog of mouse BMP-2-inducible kinase. Bone morphogenic proteins (BMPs) play a key role in skeletal development and patterning. Expression of the mouse gene is increased during BMP-2 induced differentiation and the gene product is a putative serine/threonine protein kinase containing a nuclear localization signal. Therefore, the protein encoded by this human homolog is thought to be a protein kinase with a putative regulatory role in attenuating the program of osteoblast differentiation.

# **BIKE Antibody (C-term) Blocking Peptide - References**

Arikawa, T., et al., J. Cell. Physiol. 200(3):400-406 (2004).Kearns, A.E., et al., J. Biol. Chem. 276(45):42213-42218 (2001).Hoffmann, A., et al., Crit. Rev. Eukaryot. Gene Expr. 11 (1-3), 23-45 (2001) (): ().