

EphB6 Antibody (N-term E-121) Blocking Peptide Synthetic peptide Catalog # BP7127a

Specification

EphB6 Antibody (N-term E-121) Blocking Peptide - Product Information

Primary Accession

<u>015197</u>

EphB6 Antibody (N-term E-121) Blocking Peptide - Additional Information

Gene ID 2051

Other Names Ephrin type-B receptor 6, HEP, Tyrosine-protein kinase-defective receptor EPH-6, EPHB6

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP7127a was selected from the N-terminal region with glutamic acid at position 121 of human EphB6. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions This product is for research use only. Not for use in diagnostic or therapeutic procedures.

EphB6 Antibody (N-term E-121) Blocking Peptide - Protein Information

Name EPHB6

Function

Kinase-defective receptor for members of the ephrin-B family. Binds to ephrin-B1 and ephrin-B2. Modulates cell adhesion and migration by exerting both positive and negative effects upon stimulation with ephrin-B2. Inhibits JNK activation, T-cell receptor-induced IL-2 secretion and CD25 expression upon stimulation with ephrin-B2.

Cellular Location

Membrane; Single-pass type I membrane protein.

Tissue Location

Expressed in brain. Expressed in non invasive breast carcinoma cell lines (at protein level). Strong expression in brain and pancreas, and weak expression in other tissues, such as heart, placenta, lung, liver, skeletal muscle and kidney. Expressed in breast non invasive tumors but not in



metastatic lesions. Isoform 3 is expressed in cell lines of glioblastomas, anaplastic astrocytomas, gliosarcomas and astrocytomas. Isoform 3 is not detected in normal tissues.

EphB6 Antibody (N-term E-121) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

Blocking Peptides

EphB6 Antibody (N-term E-121) Blocking Peptide - Images

EphB6 Antibody (N-term E-121) Blocking Peptide - Background

Ephrin receptors and their ligands, the ephrins, mediate numerous developmental processes, particularly in the nervous system. Based on their structures and sequence relationships, ephrins are divided into the ephrin-A (EFNA) class, which are anchored to the membrane by a glycosylphosphatidylinositol linkage, and the ephrin-B (EFNB) class, which are transmembrane proteins. The Eph family of receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. Ephrin receptor make up the largest subgroup of the receptor tyrosine kinase (RTK) family. The ephrin receptor encoded by this gene lacks the kinase activity of most receptor tyrosine kinases and binds to ephrin-B ligands.

EphB6 Antibody (N-term E-121) Blocking Peptide - References

Hafner, C., et al., Int. J. Oncol. 23(6):1553-1559 (2003).Freywald, A., et al., J. Biol. Chem. 278(12):10150-10156 (2003).Luo, H., et al., J. Clin. Invest. 110(8):1141-1150 (2002).Wilkinson, D.G., Pharmacol. Ther. 2(3):155-164 (2001).Luo, H., et al., J. Immunol. 167(3):1362-1370 (2001).