

EIF4A1 Antibody (N-term) Blocking Peptide
Synthetic peptide
Catalog # BP7262a**Specification**

EIF4A1 Antibody (N-term) Blocking Peptide - Product InformationPrimary Accession [P60842](#)**EIF4A1 Antibody (N-term) Blocking Peptide - Additional Information****Gene ID** 1973**Other Names**

Eukaryotic initiation factor 4A-I, eIF-4A-I, eIF4A-I, ATP-dependent RNA helicase eIF4A-1, EIF4A1, DDX2A, EIF4A

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP7262a](/product/products/AP7262a) was selected from the N-term region of human EIF4A1. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

EIF4A1 Antibody (N-term) Blocking Peptide - Protein Information**Name** EIF4A1**Synonyms** DDX2A, EIF4A**Function**

ATP-dependent RNA helicase which is a subunit of the eIF4F complex involved in cap recognition and is required for mRNA binding to ribosome. In the current model of translation initiation, eIF4A unwinds RNA secondary structures in the 5'-UTR of mRNAs which is necessary to allow efficient binding of the small ribosomal subunit, and subsequent scanning for the initiator codon.

EIF4A1 Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

EIF4A1 Antibody (N-term) Blocking Peptide - Images

EIF4A1 Antibody (N-term) Blocking Peptide - Background

EIF4A1 is an ATP-dependent RNA helicase which is a subunit of the eIF4F complex involved in cap recognition and is required for mRNA binding to ribosome. In the current model of translation initiation, eIF4A unwinds RNA secondary structures in the 5'-UTR of mRNAs which is necessary to allow efficient binding of the small ribosomal subunit, and subsequent scanning for the initiator codon.

EIF4A1 Antibody (N-term) Blocking Peptide - References

Suzuki,C., Proc. Natl. Acad. Sci. U.S.A. 105 (9), 3274-3279 (2008)Kim,W.J., EMBO J. 26 (24), 5020-5032 (2007)