

**LINGO1 Antibody (C-term) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP7284b****Specification**

---

**LINGO1 Antibody (C-term) Blocking Peptide - Product Information**Primary Accession [Q96FE5](#)**LINGO1 Antibody (C-term) Blocking Peptide - Additional Information**

Gene ID 84894

**Other Names**

Leucine-rich repeat and immunoglobulin-like domain-containing nogo receptor-interacting protein 1, Leucine-rich repeat and immunoglobulin domain-containing protein 1, Leucine-rich repeat neuronal protein 1, Leucine-rich repeat neuronal protein 6A, LINGO1, LERN1, LRRN6A

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP7284b](/products/AP7284b) was selected from the C-term region of human LINGO1. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**LINGO1 Antibody (C-term) Blocking Peptide - Protein Information**

Name LINGO1

Synonyms LERN1, LRRN6A

**Function**

Functional component of the Nogo receptor signaling complex (RTN4R/NGFR) in RhoA activation responsible for some inhibition of axonal regeneration by myelin-associated factors (PubMed: [14966521](http://www.uniprot.org/citations/14966521), PubMed: [15694321](http://www.uniprot.org/citations/15694321)). Is also an important negative regulator of oligodendrocyte differentiation and axonal myelination (PubMed: [15895088](http://www.uniprot.org/citations/15895088)). Acts in conjunction with RTN4 and RTN4R in regulating neuronal precursor cell motility during cortical development (By similarity).

**Cellular Location**

Cell membrane {ECO:0000250|UniProtKB:Q9D1T0}; Single-pass type I membrane protein {ECO:0000250|UniProtKB:Q9D1T0}

**Tissue Location**

Expressed exclusively in the central nervous system. Highest level in the in amygdala, hippocampus, thalamus and cerebral cortex. In the rest of the brain a basal expression seems to be always present. Up-regulated in substantia nigra neurons from Parkinson disease patients.

**LINGO1 Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**LINGO1 Antibody (C-term) Blocking Peptide - Images****LINGO1 Antibody (C-term) Blocking Peptide - Background**

LINGO1 is a functional component of the Nogo receptor signaling complex(RTN4R/NGFR) in RhoA activation responsible for some inhibition of axonal regeneration by myelin-associated factors. It is also an important negative regulator of oligodendrocyte differentiation and axonal myelination.

**LINGO1 Antibody (C-term) Blocking Peptide - References**

Inoue,H., Proc. Natl. Acad. Sci. U.S.A. 104 (36), 14430-14435 (2007)Satoh,J., Neuropathol. Appl. Neurobiol. 33 (1), 99-107 (2007)Mosyak,L., J. Biol. Chem. 281 (47), 36378-36390 (2006)Mi,S., Nat. Neurosci. 7 (3), 221-228 (2004)