

CYK18 Antibody (C-term) Blocking Peptide

Synthetic peptide Catalog # BP7376b

Specification

CYK18 Antibody (C-term) Blocking Peptide - Product Information

Primary Accession

P05783

CYK18 Antibody (C-term) Blocking Peptide - Additional Information

Gene ID 3875

Other Names

Keratin, type I cytoskeletal 18, Cell proliferation-inducing gene 46 protein, Cytokeratin-18, CK-18, Keratin-18, K18, KRT18, CYK18

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP7376b was selected from the C-term region of human CYK18. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

CYK18 Antibody (C-term) Blocking Peptide - Protein Information

Name KRT18

Synonyms CYK18

Function

Involved in the uptake of thrombin-antithrombin complexes by hepatic cells (By similarity). When phosphorylated, plays a role in filament reorganization. Involved in the delivery of mutated CFTR to the plasma membrane. Together with KRT8, is involved in interleukin-6 (IL-6)-mediated barrier protection.

Cellular Location

Nucleus matrix {ECO:0000250|UniProtKB:Q5BJY9}. Cytoplasm, perinuclear region. Nucleus, nucleolus. Cytoplasm {ECO:0000250|UniProtKB:Q5BJY9}



Tissue Location

Expressed in colon, placenta, liver and very weakly in exocervix. Increased expression observed in lymph nodes of breast carcinoma.

CYK18 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides

CYK18 Antibody (C-term) Blocking Peptide - Images

CYK18 Antibody (C-term) Blocking Peptide - Background

KRT18 is the type I intermediate filament chain keratin 18. Keratin 18, together with its filament partner keratin 8, are perhaps the most commonly found members of the intermediate filament family. They are expressed in single layer epithelial tissues of the body. Mutations in its gene have been linked to cryptogenic cirrhosis.

CYK18 Antibody (C-term) Blocking Peptide - References

Zhang, Q., Clin. Cancer Res. 15 (10), 3557-3567 (2009) Kruse, R., Folia Histochem. Cytobiol. 47 (1), 127-130 (2009) Toivola, D.M., Hepatology 40 (2), 459-466 (2004)