

**STK23 (MSSK1) Antibody (Center) Blocking peptide**  
**Synthetic peptide**  
**Catalog # BP7544c****Specification**

---

**STK23 (MSSK1) Antibody (Center) Blocking peptide - Product Information**

Primary Accession [O9UPE1](#)  
Other Accession [NP\\_001164232](#)

**STK23 (MSSK1) Antibody (Center) Blocking peptide - Additional Information**

**Gene ID** 26576

**Other Names**

SRSF protein kinase 3, Muscle-specific serine kinase 1, MSSK-1, Serine/arginine-rich protein-specific kinase 3, SR-protein-specific kinase 3, Serine/threonine-protein kinase 23, SRPK3, MSSK1, STK23

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP7544c](/product/products/AP7544c) was selected from the Center region of human MSSK1 . A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**STK23 (MSSK1) Antibody (Center) Blocking peptide - Protein Information**

**Name** SRPK3

**Synonyms** MSSK1, STK23

**Function**

Serine/arginine-rich protein-specific kinase which specifically phosphorylates its substrates at serine residues located in regions rich in arginine/serine dipeptides, known as RS domains. Phosphorylates the SR splicing factor SRSF1 and the lamin-B receptor (LBR) in vitro. Required for normal muscle development (By similarity).

**Tissue Location**

Exclusively expressed in skeletal and heart muscle.

**STK23 (MSSK1) Antibody (Center) Blocking peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**STK23 (MSSK1) Antibody (Center) Blocking peptide - Images****STK23 (MSSK1) Antibody (Center) Blocking peptide - Background**

Protein kinases are enzymes that transfer a phosphate group from a phosphate donor, generally the  $\gamma$  phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this basic mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells, regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement and cell movement, apoptosis, and differentiation. With more than 500 gene products, the protein kinase family is one of the largest families of proteins in eukaryotes. The family has been classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or serine/threonine (STK) kinase catalytic domains. The STE group (homologs of yeast Sterile 7, 11, 20 kinases) consists of 50 kinases related to the mitogen-activated protein kinase (MAPK) cascade families (Ste7/MAP2K, Ste11/MAP3K, and Ste20/MAP4K). MAP kinase cascades, consisting of a MAPK and one or more upstream regulatory kinases (MAPKKs) have been best characterized in the yeast pheromone response pathway. Pheromones bind to Ste cell surface receptors and activate yeast MAPK pathway.

**STK23 (MSSK1) Antibody (Center) Blocking peptide - References**

Brenner, V., et al., Genomics 44(1):8-14 (1997).