

PFTK1 Antibody (N-term) Blocking Peptide
Synthetic peptide
Catalog # BP7550a**Specification**

PFTK1 Antibody (N-term) Blocking Peptide - Product Information

Primary Accession [O94921](#)
Other Accession [NP_036527](#)

PFTK1 Antibody (N-term) Blocking Peptide - Additional Information

Gene ID 5218

Other Names

Cyclin-dependent kinase 14, Cell division protein kinase 14, Serine/threonine-protein kinase PFTK1, hPFTK1, CDK14, KIAA0834, PFTK1

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP7550a](#) was selected from the N-term region of human PFTK1. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

PFTK1 Antibody (N-term) Blocking Peptide - Protein Information

Name CDK14

Synonyms KIAA0834, PFTK1

Function

Serine/threonine-protein kinase involved in the control of the eukaryotic cell cycle, whose activity is controlled by an associated cyclin. Acts as a cell-cycle regulator of Wnt signaling pathway during G2/M phase by mediating the phosphorylation of LRP6 at 'Ser-1490', leading to the activation of the Wnt signaling pathway. Acts as a regulator of cell cycle progression and cell proliferation via its interaction with CCDN3. Phosphorylates RB1 in vitro, however the relevance of such result remains to be confirmed in vivo. May also play a role in meiosis, neuron differentiation and may indirectly act as a negative regulator of insulin-responsive glucose transport.

Cellular Location

Cell membrane; Peripheral membrane protein. Cytoplasm. Nucleus. Note=Recruited to the cell membrane by CCNY

Tissue Location

Highly expressed in brain, pancreas, kidney, heart, testis and ovary. Also detected at lower levels in other tissues except in spleen and thymus where expression is barely detected

PFTK1 Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

PFTK1 Antibody (N-term) Blocking Peptide - Images**PFTK1 Antibody (N-term) Blocking Peptide - Background**

PFTK1, a member of the CDC2/CDKX subfamily of Ser/Thr protein kinases, may play a role in meiosis as well as in neuron differentiation and/or function. It is highly expressed in brain, pancreas, kidney, heart, testis and ovary, and also detected at lower levels in other tissues except in spleen and thymus where expression is minimal.

PFTK1 Antibody (N-term) Blocking Peptide - References

Yang, T., et al., Gene 267(2):165-172 (2001). Nagase, T., et al., DNA Res. 5(6):355-364 (1998).