

**HEPTP (PTPN7) Antibody (N-term M1) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP7599b****Specification**

---

**HEPTP (PTPN7) Antibody (N-term M1) Blocking Peptide - Product Information**Primary Accession [P35236](#)**HEPTP (PTPN7) Antibody (N-term M1) Blocking Peptide - Additional Information****Gene ID** 5778**Other Names**

Tyrosine-protein phosphatase non-receptor type 7, Hematopoietic protein-tyrosine phosphatase, HEPTP, Protein-tyrosine phosphatase LC-PTP, PTPN7

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP7599b](/products/AP7599b) was selected from the PTPN7 region of human HEPTP (PTPN7). A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**HEPTP (PTPN7) Antibody (N-term M1) Blocking Peptide - Protein Information****Name** PTPN7**Function**

Protein phosphatase that acts preferentially on tyrosine- phosphorylated MAPK1. Plays a role in the regulation of T and B- lymphocyte development and signal transduction.

**Cellular Location**

Cytoplasm. Cytoplasm, cytoskeleton

**Tissue Location**

Expressed exclusively in thymus and spleen.



## **HEPTP (PTPN7) Antibody (N-term M1) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

## **HEPTP (PTPN7) Antibody (N-term M1) Blocking Peptide - Images**

## **HEPTP (PTPN7) Antibody (N-term M1) Blocking Peptide - Background**

PTPN7 is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. PTPN7 is preferentially expressed in a variety of hematopoietic cells, and is an early response gene in lymphokine stimulated cells. The noncatalytic N-terminus of this PTP can interact with MAP kinases and suppress the MAP kinase activities. This PTP has been shown to be involved in the regulation of T cell antigen receptor (TCR) signaling, which is thought to function through dephosphorylating the molecules related to MAP kinase pathway.

## **HEPTP (PTPN7) Antibody (N-term M1) Blocking Peptide - References**

Eswaran,J., Biochem. J. 395 (3), 483-491 (2006)Mustelin,T., J. Mol. Biol. 354 (1), 150-163 (2005)Pettiford,S.M., Leukemia 17 (2), 366-378 (2003)Kosaki,K., J. Clin. Endocrinol. Metab. 87 (8), 3529-3533 (2002)