

**MGRN1 Antibody (C-term) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP7634b****Specification**

---

**MGRN1 Antibody (C-term) Blocking Peptide - Product Information**Primary Accession [O60291](#)**MGRN1 Antibody (C-term) Blocking Peptide - Additional Information****Gene ID** 23295**Other Names**

E3 ubiquitin-protein ligase MGRN1, 632-, Mahogunin RING finger protein 1, RING finger protein 156, MGRN1, KIAA0544, RNF156

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP7634b](/products/AP7634b) was selected from the C-term region of human MGRN1. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**MGRN1 Antibody (C-term) Blocking Peptide - Protein Information****Name** MGRN1**Synonyms** KIAA0544, RNF156**Function**

E3 ubiquitin-protein ligase. Mediates monoubiquitination at multiple sites of TSG101 in the presence of UBE2D1, but not of UBE2G1, nor UBE2H. Plays a role in the regulation of endosome-to-lysosome trafficking. Impairs MC1R- and MC4R-signaling by competing with GNAS-binding to MCRs and inhibiting agonist-induced cAMP production. Does not inhibit ADRB2-signaling. Does not promote MC1R ubiquitination. Acts also as a negative regulator of hedgehog signaling (By similarity).

**Cellular Location**

Early endosome. Note=The endosomal localization is dependent on the interaction with TSG101

[Isoform 2]: Cytoplasm, cytosol. Nucleus. Note=Translocation from the cytosol to the nucleus is seen only in the presence of MC1R and MC4R, but not TBXA2R. Excluded from nucleoli [Isoform 4]: Cytoplasm, cytosol. Cell membrane.

### **MGRN1 Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

### **MGRN1 Antibody (C-term) Blocking Peptide - Images**

### **MGRN1 Antibody (C-term) Blocking Peptide - Background**

Mahogunin (MGRN1) is a C3HC4 RING-containing protein with E3 ubiquitin ligase activity in vitro.

### **MGRN1 Antibody (C-term) Blocking Peptide - References**

Chakrabarti,O., Cell 137 (6), 1136-1147 (2009)Kim,B.Y., Mol. Biol. Cell 18 (4), 1129-1142 (2007)