

Catalog # BP7735a

Runx2 Antibody (N-term) Blocking Peptide Synthetic peptide

## Specification

# Runx2 Antibody (N-term) Blocking Peptide - Product Information

Primary Accession

<u>Q13950</u>

### Runx2 Antibody (N-term) Blocking Peptide - Additional Information

Gene ID 860

#### **Other Names**

Runt-related transcription factor 2, Acute myeloid leukemia 3 protein, Core-binding factor subunit alpha-1, CBF-alpha-1, Oncogene AML-3, Osteoblast-specific transcription factor 2, OSF-2, Polyomavirus enhancer-binding protein 2 alpha A subunit, PEA2-alpha A, PEBP2-alpha A, SL3-3 enhancer factor 1 alpha A subunit, SL3/AKV core-binding factor alpha A subunit, RUNX2, AML3, CBFA1, OSF2, PEBP2A

#### Target/Specificity

The synthetic peptide sequence used to generate the antibody <a href=/products/AP7735a>AP7735a</a> was selected from the N-term region of human Runx2. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage** Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

### Runx2 Antibody (N-term) Blocking Peptide - Protein Information

Name RUNX2

Synonyms AML3, CBFA1, OSF2, PEBP2A

Function

Transcription factor involved in osteoblastic differentiation and skeletal morphogenesis (PubMed:<a href="http://www.uniprot.org/citations/28505335" target="\_blank">28505335</a>, PubMed:<a href="http://www.uniprot.org/citations/28738062" target="\_blank">28738062</a>, PubMed:<a href="http://www.uniprot.org/citations/28703881" target="\_blank">28703881</a>, PubMed:<a href="http://www.uniprot.org/citations/28703881" target="\_blank">28703881</a>, PubMed:<a href="http://www.uniprot.org/citations/28703881" target="\_blank">28703881</a>, PubMed:<a href="http://www.uniprot.org/citations/28703881" target="\_blank">28703881</a>, PubMed:<a href="http://www.uniprot.org/citations/28703881" target="\_blank">28703881</a>). Essential for the maturation of osteoblasts and both intramembranous and endochondral ossification. CBF binds to the core site, 5'-PYGPYGGT-3', of a number of enhancers and promoters,



including murine leukemia virus, polyomavirus enhancer, T-cell receptor enhancers, osteocalcin, osteopontin, bone sialoprotein, alpha 1(I) collagen, LCK, IL-3 and GM-CSF promoters. In osteoblasts, supports transcription activation: synergizes with SPEN/MINT to enhance FGFR2-mediated activation of the osteocalcin FGF-responsive element (OCFRE) (By similarity). Inhibits KAT6B-dependent transcriptional activation.

Cellular Location Nucleus. Cytoplasm {ECO:0000250|UniProtKB:Q08775}

**Tissue Location** Specifically expressed in osteoblasts.

## Runx2 Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

<u>Blocking Peptides</u>

#### Runx2 Antibody (N-term) Blocking Peptide - Images

### Runx2 Antibody (N-term) Blocking Peptide - Background

Runx2 is a member of the RUNX family of transcription factors. It is a nuclear protein with an Runt DNA-binding domain. This protein is essential for osteoblastic differentiation and skeletal morphogenesis and acts as a scaffold for nucleic acids and regulatory factors involved in skeletal gene expression. It can bind DNA both as a monomer or, with more affinity, as a subunit of a heterodimeric complex. Mutations in the Runx2 gene have been associated with the bone development disorder cleidocranial dysplasia (CCD).

### Runx2 Antibody (N-term) Blocking Peptide - References

Rich,J.T., Biochem. Biophys. Res. Commun. 372 (1), 230-235 (2008)Ermakov,S., Ann. Hum. Genet. 72 (PT 4), 510-518 (2008)Endo,T., J. Clin. Endocrinol. Metab. 93 (6), 2409-2412 (2008)