

**APS Antibody (Center) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP7747c****Specification**

---

**APS Antibody (Center) Blocking Peptide - Product Information**Primary Accession [O14492](#)**APS Antibody (Center) Blocking Peptide - Additional Information****Gene ID** 10603**Other Names**

SH2B adapter protein 2, Adapter protein with pleckstrin homology and Src homology 2 domains, SH2 and PH domain-containing adapter protein APS, SH2B2, APS

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP7747c](/products/AP7747c) was selected from the Center region of human APS. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**APS Antibody (Center) Blocking Peptide - Protein Information****Name** SH2B2**Synonyms** APS**Function**

Adapter protein for several members of the tyrosine kinase receptor family. Involved in multiple signaling pathways. May be involved in coupling from immunoreceptor to Ras signaling. Acts as a negative regulator of cytokine signaling in collaboration with CBL. Binds to EPOR and suppresses EPO-induced STAT5 activation, possibly through a masking effect on STAT5 docking sites in EPOR. Suppresses PDGF-induced mitogenesis. May induce cytoskeletal reorganization via interaction with VAV3.

**Cellular Location**

Cytoplasm. Cell membrane. Note=Cytoplasmic before PDGF stimulation. After PDGF stimulation,

localized at the cell membrane and peripheral region

**Tissue Location**

Expressed in spleen, prostate, testis, uterus, small intestine and skeletal muscle. Among hematopoietic cell lines, expressed exclusively in B-cells. Not expressed in most tumor cell lines.

**APS Antibody (Center) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**APS Antibody (Center) Blocking Peptide - Images****APS Antibody (Center) Blocking Peptide - Background**

APS is expressed in B lymphocytes and contains pleckstrin homology and src homology 2 (SH2) domains. In Burkitt lymphoma cell lines, it is tyrosine phosphorylated in response to B cell receptor stimulation. Because it binds Shc independent of stimulation and Grb2 after stimulation, it appears to play a role in signal transduction from the receptor to Shc/Grb2.

**APS Antibody (Center) Blocking Peptide - References**

Li,M., Endocrinology 148 (4), 1615-1621 (2007)Katsanakis,K.D., J. Biol. Chem. 280 (45), 37827-37832 (2005)Iseki,M., Biochem. Biophys. Res. Commun. 330 (3), 1005-1013 (2005)