

MORF/MYST4 Antibody (N-term) Blocking Peptide
Synthetic peptide
Catalog # BP7828a**Specification**

MORF/MYST4 Antibody (N-term) Blocking Peptide - Product InformationPrimary Accession [Q8WYB5](#)**MORF/MYST4 Antibody (N-term) Blocking Peptide - Additional Information****Gene ID** 23522**Other Names**

Histone acetyltransferase KAT6B, Histone acetyltransferase MOZ2, MOZ, YBF2/SAS3, SAS2 and TIP60 protein 4, MYST-4, Monocytic leukemia zinc finger protein-related factor, KAT6B, KIAA0383, MORF, MOZ2, MYST4

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP7828a](/product/products/AP7828a) was selected from the N-term region of human MORF/MYST4. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

MORF/MYST4 Antibody (N-term) Blocking Peptide - Protein Information**Name** KAT6B**Synonyms** KIAA0383, MORF, MOZ2, MYST4**Function**

Histone acetyltransferase which may be involved in both positive and negative regulation of transcription. Required for RUNX2- dependent transcriptional activation. May be involved in cerebral cortex development. Component of the MOZ/MORF complex which has a histone H3 acetyltransferase activity.

Cellular Location

Nucleus.

Tissue Location

Ubiquitously expressed, with high levels in heart, pancreas, testis and ovary.

MORF/MYST4 Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

MORF/MYST4 Antibody (N-term) Blocking Peptide - Images**MORF/MYST4 Antibody (N-term) Blocking Peptide - Background**

MORF/MYST4 is a histone acetyltransferase which may be involved in both positive and negative regulation of transcription. It is required for RUNX2-dependent transcriptional activation and may be involved in cerebral cortex development.

MORF/MYST4 Antibody (N-term) Blocking Peptide - References

Pena,A.N., Ann. N. Y. Acad. Sci. 1100, 299-305 (2007)Olsen,J.V., Cell 127 (3), 635-648 (2006)Liu,C., Cytokine 27 (4-5), 93-100 (2004)