

AAK1 Antibody (C-term) Blocking Peptide Synthetic peptide Catalog # BP7861b

Specification

AAK1 Antibody (C-term) Blocking Peptide - Product Information

Primary Accession

<u>Q2M2I8</u>

AAK1 Antibody (C-term) Blocking Peptide - Additional Information

Gene ID 22848

Other Names AP2-associated protein kinase 1, Adaptor-associated kinase 1, AAK1, KIAA1048

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP7861b was selected from the C-term region of human AAK1. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

AAK1 Antibody (C-term) Blocking Peptide - Protein Information

Name AAK1

Synonyms KIAA1048

Function

Regulates clathrin-mediated endocytosis by phosphorylating the AP2M1/mu2 subunit of the adaptor protein complex 2 (AP-2) which ensures high affinity binding of AP-2 to cargo membrane proteins during the initial stages of endocytosis (PubMed:<a

```
href="http://www.uniprot.org/citations/17494869" target="_blank">17494869</a>, PubMed:<a href="http://www.uniprot.org/citations/11877457" target="_blank">11877457</a>, PubMed:<a href="http://www.uniprot.org/citations/11877461" target="_blank">11877461</a>, PubMed:<a href="http://www.uniprot.org/citations/12952931" target="_blank">12952931</a>, PubMed:<a href="http://www.uniprot.org/citations/12952931" target="_blank">14617351</a>, PubMed:<a href="http://www.uniprot.org/citations/14617351" target="_blank">14617351</a>, PubMed:<a href="http://www.uniprot.org/citations/14617351" target="_blank">14617351</a>, PubMed:<a href="http://www.uniprot.org/citations/14617351" target="_blank">25653444</a>). Isoform 1 and isoform 2 display similar levels of kinase activity towards AP2M1 (PubMed:<a
```



href="http://www.uniprot.org/citations/17494869" target="_blank">17494869).
Preferentially, may phosphorylate substrates on threonine residues (PubMed:11877457, PubMed:18657069). Regulates
phosphorylation of other AP-2 subunits as well as AP-2 localization and AP-2-mediated
internalization of ligand complexes (PubMed:<a href="http://www.uniprot.org/citations/12952931"
target="_blank">12952931). Phosphorylates NUMB and regulates its cellular localization,
promoting NUMB localization to endosomes (PubMed:18657069). Binds to and
stabilizes the activated form of NOTCH1, increases its localization in endosomes and regulates its
transcriptional activity (PubMed:<a href="http://www.uniprot.org/citations/21464124"
target="_blank">21464124).

Cellular Location

Cell membrane {ECO:0000250|UniProtKB:F1MH24}; Peripheral membrane protein {ECO:0000250|UniProtKB:F1MH24}. Membrane, clathrin-coated pit. Presynapse {ECO:0000250|UniProtKB:P0C1X8}. Note=Active when found in clathrin- coated pits at the plasma membrane. In neuronal cells, enriched at presynaptic terminals. In non-neuronal cells, enriched at leading edge of migrating cells. {ECO:0000250|UniProtKB:P0C1X8}

Tissue Location

Detected in brain, heart and liver. Isoform 1 is the predominant isoform in brain.

AAK1 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

<u>Blocking Peptides</u>

AAK1 Antibody (C-term) Blocking Peptide - Images

AAK1 Antibody (C-term) Blocking Peptide - Background

Adaptor-related protein complex 2 (AP-2 complexes) functions during receptor-mediated endocytosis to trigger clathrin assembly, interact with membrane-bound receptors, and recruit encodytic accessory factors. AAK1 is a member of the SNF1 subfamily of Ser/Thr protein kinases. The protein interacts with and phosphorylates a subunit of the AP-2 complex, which promotes binding of AP-2 to sorting signals found in membrane-bound receptors and subsequent receptor endocytosis. Its kinase activity is stimulated by clathrin.

AAK1 Antibody (C-term) Blocking Peptide - References

Henderson, D.M., Mol. Biol. Cell 18 (7), 2698-2706 (2007) Takahashi, T., Cancer Res. 66 (24), 11932-11937 (2006) Schmid, E.M., PLoS Biol. 4 (9), E262 (2006)