

**MLK1 Antibody (C-term) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP7919a****Specification**

---

**MLK1 Antibody (C-term) Blocking Peptide - Product Information**Primary Accession [P80192](#)**MLK1 Antibody (C-term) Blocking Peptide - Additional Information****Gene ID** 4293**Other Names**

Mitogen-activated protein kinase kinase kinase 9, Mixed lineage kinase 1, MAP3K9, MLK1, PRKE1

**Target/Specificity**

The synthetic peptide sequence is selected from aa 1046~1066 of human MLK1.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**MLK1 Antibody (C-term) Blocking Peptide - Protein Information****Name** MAP3K9**Synonyms** MLK1, PRKE1**Function**

Serine/threonine kinase which acts as an essential component of the MAP kinase signal transduction pathway. Plays an important role in the cascades of cellular responses evoked by changes in the environment. Once activated, acts as an upstream activator of the MKK/JNK signal transduction cascade through the phosphorylation of MAP2K4/MKK4 and MAP2K7/MKK7 which in turn activate the JNKs. The MKK/JNK signaling pathway regulates stress response via activator protein-1 (JUN) and GATA4 transcription factors. Also plays a role in mitochondrial death signaling pathway, including the release cytochrome c, leading to apoptosis.

**Tissue Location**

Expressed in epithelial tumor cell lines of colonic, breast and esophageal origin.

## **MLK1 Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

## **MLK1 Antibody (C-term) Blocking Peptide - Images**

## **MLK1 Antibody (C-term) Blocking Peptide - Background**

Protein kinases are enzymes that transfer a phosphate group from a phosphate donor, generally the  $\gamma$  phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this basic mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells, regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement and cell movement, apoptosis, and differentiation. With more than 500 gene products, the protein kinase family is one of the largest families of proteins in eukaryotes. The family has been classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or serine/threonine (STK) kinase catalytic domains. The tyrosine-like kinase (TLK) group consists of 40 tyrosine and serine-threonine kinases such as MLK (mixed-lineage kinase), LSK (LIMK/TESK), IRAK (interleukin-1 receptor-associated kinase), Raf, RIPK (receptor-interacting protein kinase), and STRK (activin and TGF-beta receptors) families.

## **MLK1 Antibody (C-term) Blocking Peptide - References**

Blume-Jensen P, et al. Nature 2001. 411: 355. Cantrell D, J. Cell Sci. 2001. 114: 1439. Jhang S  
Oncogene 2000. 19: 5590. Manning G, et al. Science 2002. 298: 1912. Moller, D, et al. Am. J. Physiol.  
1994. 266: C351-C359. Robertson, S. et al. Trends Genet. 2000. 16: 368. Robinson D, et al.  
Oncogene 2000. 19: 5548. Van der Ven, P, et al. Hum. Molec. Genet. 1993. 2: 1889. Vanhaesebroeck,  
B, et al. Biochem. J. 2000. 346: 561. Van Weering D, et al. Recent Results Cancer Res. 1998. 154:  
271.