

KHS2 Antibody (C-term) Blocking Peptide
Synthetic peptide
Catalog # BP7972b**Specification**

KHS2 Antibody (C-term) Blocking Peptide - Product InformationPrimary Accession
Other Accession[Q8IVH8](#)
[NP_003609](#)**KHS2 Antibody (C-term) Blocking Peptide - Additional Information****Gene ID** 8491**Other Names**

Mitogen-activated protein kinase kinase kinase 3, Germinal center kinase-related protein kinase, GLK, MAPK/ERK kinase kinase kinase 3, MEK kinase kinase 3, MEKKK 3, MAP4K3, RAB8IPL1

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP7972b](/product/products/AP7972b) was selected from the C-term region of human KHS2 . A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

KHS2 Antibody (C-term) Blocking Peptide - Protein Information**Name** MAP4K3**Synonyms** RAB8IPL1**Function**

May play a role in the response to environmental stress. Appears to act upstream of the JUN N-terminal pathway.

Tissue Location

Ubiquitously expressed in all tissues examined, with high levels in heart, brain, placenta, skeletal muscle, kidney and pancreas and lower levels in lung and liver

KHS2 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

KHS2 Antibody (C-term) Blocking Peptide - Images

KHS2 Antibody (C-term) Blocking Peptide - Background

KHS2, a member of the Ste20 subfamily of Ser/Thr protein kinases, features the characteristic N-terminal catalytic domain and C-terminal regulatory domain of the Ste20 subfamily. The kinase activity of the encoded protein can be stimulated by UV radiation and tumor necrosis factor- α . The protein specifically activates the c-Jun N-terminal kinase (JNK) signaling pathway. Evidence suggests that it functions upstream of mitogen-activated protein kinase kinase kinase 1 (MEKK1). This gene previously was referred to as RAB8-interacting protein-like 1 (RAB8IPL1), but it has been renamed mitogen-activated protein kinase kinase kinase kinase 3 (MAP4K3).

KHS2 Antibody (C-term) Blocking Peptide - References

Diener, K., et al., Proc. Natl. Acad. Sci. U.S.A. 94(18):9687-9692 (1997).