

PIP5KI gamma (PIP5K1G) Antibody (C-term) Blocking peptide
Synthetic peptide
Catalog # BP8039b

Specification

PIP5KI gamma (PIP5K1G) Antibody (C-term) Blocking peptide - Product Information

Primary Accession [O60331](#)

PIP5KI gamma (PIP5K1G) Antibody (C-term) Blocking peptide - Additional Information

Gene ID 23396

Other Names

Phosphatidylinositol 4-phosphate 5-kinase type-1 gamma, PIP5K1-gamma, PtdIns(4)P-5-kinase 1 gamma, Phosphatidylinositol 4-phosphate 5-kinase type I gamma, PIP5KIgamma, PIP5K1C, KIAA0589

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP8039b](/product/products/AP8039b) was selected from the C-term region of human PIP5K1G . A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

PIP5KI gamma (PIP5K1G) Antibody (C-term) Blocking peptide - Protein Information

Name PIP5K1C ([HGNC:8996](#))

Synonyms KIAA0589

Function

Catalyzes the phosphorylation of phosphatidylinositol 4- phosphate (PtdIns(4)P/PI4P) to form phosphatidylinositol 4,5- biphosphate (PtdIns(4,5)P2/PIP2), a lipid second messenger that regulates several cellular processes such as signal transduction, vesicle trafficking, actin cytoskeleton dynamics, cell adhesion, and cell motility (PubMed:[12422219](http://www.uniprot.org/citations/12422219), PubMed:[22942276](http://www.uniprot.org/citations/22942276)). PtdIns(4,5)P2 can directly act as a second messenger or can be utilized as a precursor to generate other second messengers: inositol 1,4,5-trisphosphate (IP3), diacylglycerol (DAG) or

phosphatidylinositol-3,4,5-trisphosphate (PtdIns(3,4,5)P3/PIP3) (Probable). PIP5K1A-mediated phosphorylation of PtdIns(4)P is the predominant pathway for PtdIns(4,5)P2 synthesis (By similarity). Together with PIP5K1A, is required for phagocytosis, both enzymes regulating different types of actin remodeling at sequential steps (By similarity). Promotes particle attachment by generating the pool of PtdIns(4,5)P2 that induces controlled actin depolymerization to facilitate Fc-gamma-R clustering. Mediates RAC1-dependent reorganization of actin filaments. Required for synaptic vesicle transport (By similarity). Controls the plasma membrane pool of PtdIns(4,5)P2 implicated in synaptic vesicle endocytosis and exocytosis (PubMed:12847086). Plays a role in endocytosis mediated by clathrin and AP-2 (adaptor protein complex 2) (PubMed:12847086). Required for clathrin-coated pits assembly at the synapse (PubMed:17261850). Participates in cell junction assembly (PubMed:17261850). Modulates adherens junctions formation by facilitating CDH1/cadherin trafficking (PubMed:17261850). Required for focal adhesion dynamics. Modulates the targeting of talins (TLN1 and TLN2) to the plasma membrane and their efficient assembly into focal adhesions (PubMed:12422219). Regulates the interaction between talins (TLN1 and TLN2) and beta-integrins (PubMed:12422219). Required for uropodium formation and retraction of the cell rear during directed migration (By similarity). Has a role in growth factor-stimulated directional cell migration and adhesion (By similarity). Required for talin assembly into nascent adhesions forming at the leading edge toward the direction of the growth factor (PubMed:17635937). Negative regulator of T-cell activation and adhesion (By similarity). Negatively regulates integrin alpha-L/beta-2 (LFA-1) polarization and adhesion induced by T-cell receptor (By similarity). Together with PIP5K1A has a role during embryogenesis and together with PIP5K1B may have a role immediately after birth (By similarity).

Cellular Location

Cell membrane; Peripheral membrane protein; Cytoplasmic side {ECO:0000250|UniProtKB:Q5I6B8}. Endomembrane system {ECO:0000250|UniProtKB:Q5I6B8}. Cytoplasm {ECO:0000250|UniProtKB:O70161}. Cell junction, focal adhesion. Cell junction, adherens junction. Cell projection, ruffle membrane {ECO:0000250|UniProtKB:Q5I6B8}. Cell projection, phagocytic cup {ECO:0000250|UniProtKB:O70161}. Cell projection, uropodium {ECO:0000250|UniProtKB:O70161}. Note=Detected in plasma membrane invaginations. Isoform 3 is detected in intracellular vesicle-like structures

Tissue Location

[Isoform 1]: Isoform 1 is strongly expressed in brain and also detected in heart and lung [Isoform 3]: Isoform 3 is detected in large amounts in heart and large intestine, is also present in lung, pancreas and thyroid, and to a lesser extent in brain, stomach and kidney

PIP5KI gamma (PIP5K1G) Antibody (C-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

PIP5KI gamma (PIP5K1G) Antibody (C-term) Blocking peptide - Images

PIP5KI gamma (PIP5K1G) Antibody (C-term) Blocking peptide - Background

Protein kinases are enzymes that transfer a phosphate group from a phosphate donor, generally the gamma phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this basic

mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells, regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement and cell movement, apoptosis, and differentiation. With more than 500 gene products, the protein kinase family is one of the largest families of proteins in eukaryotes. The family has been classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or serine/threonine (STK) kinase catalytic domains.

PIP5KI gamma (PIP5K1G) Antibody (C-term) Blocking peptide - References

Ling, K., et al., Nature 420(6911):89-93 (2002). Di Paolo, G., et al., Nature 420(6911):85-89 (2002). Ishihara, H., et al., J. Biol. Chem. 273(15):8741-8748 (1998).