

PACSIN2 Antibody (C-term) Blocking Peptide
Synthetic peptide
Catalog # BP8088b

Specification

PACSIN2 Antibody (C-term) Blocking Peptide - Product Information

Primary Accession [Q9UNF0](#)

PACSIN2 Antibody (C-term) Blocking Peptide - Additional Information

Gene ID 11252

Other Names

Protein kinase C and casein kinase substrate in neurons protein 2, Syndapin-2, Syndapin-II, PACSIN2

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP8088b](/product/products/AP8088b) was selected from the C-term region of human PACSIN2 . A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

PACSIN2 Antibody (C-term) Blocking Peptide - Protein Information

Name PACSIN2

Function

Regulates the morphogenesis and endocytosis of caveolae (By similarity). Lipid-binding protein that is able to promote the tubulation of the phosphatidic acid-containing membranes it preferentially binds. Plays a role in intracellular vesicle-mediated transport. Involved in the endocytosis of cell-surface receptors like the EGF receptor, contributing to its internalization in the absence of EGF stimulus.

Cellular Location

Cytoplasm. Cytoplasm, cytoskeleton. Cytoplasmic vesicle membrane; Peripheral membrane protein; Cytoplasmic side. Early endosome. Recycling endosome membrane Cell projection, ruffle membrane; Peripheral membrane protein; Cytoplasmic side. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection Membrane, caveola. Note=Detected at the

neck of flask-shaped caveolae. Localization to tubular recycling endosomes probably requires interaction with MICALL1 and EHD1

Tissue Location

Widely expressed.

PACSIN2 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

PACSIN2 Antibody (C-term) Blocking Peptide - Images**PACSIN2 Antibody (C-term) Blocking Peptide - Background**

PACSIN may play a role in vesicle formation and transport. This protein homo- and hetero-aggregates with other PACSINs. It also binds dynamin 1, synaptojanin, synapsin 1 and the neural Wiskott-Aldrich syndrome protein (N-WASP). The protein exhibits a cvesicle-like cytoplasmic distribution and is ubiquitously expressed. PACSIN is phosphorylated by casein kinase 2 (CK2) and protein kinase C (PKC). The protein contains 1 FCH domain and 1 SH3 domain.

PACSIN2 Antibody (C-term) Blocking Peptide - References

Strausberg, R.L., et al., Proc. Natl. Acad. Sci. U.S.A. 99(26):16899-16903 (2002). Wiemann, S., et al., Genome Res. 11(3):422-435 (2001). Ritter, B., et al., FEBS Lett. 454(3):356-362 (1999). Dunham, I., et al., Nature 402(6761):489-495 (1999).