

DGKE Antibody (C-term) Blocking Peptide
Synthetic peptide
Catalog # BP8125b**Specification**

DGKE Antibody (C-term) Blocking Peptide - Product InformationPrimary Accession [P52429](#)**DGKE Antibody (C-term) Blocking Peptide - Additional Information****Gene ID** 8526**Other Names**

Diacylglycerol kinase epsilon, DAG kinase epsilon, Diglyceride kinase epsilon, DGK-epsilon, DGKE, DAGK5

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP8125b](/product/products/AP8125b) was selected from the C-term region of human DGKE. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

DGKE Antibody (C-term) Blocking Peptide - Protein Information**Name** DGKE**Synonyms** DAGK5**Function**

Membrane-bound diacylglycerol kinase that converts diacylglycerol/DAG into phosphatidic acid/phosphatidate/PA and regulates the respective levels of these two bioactive lipids (PubMed: [15544348](http://www.uniprot.org/citations/15544348), PubMed: [19744926](http://www.uniprot.org/citations/19744926), PubMed: [22108654](http://www.uniprot.org/citations/22108654), PubMed: [21477596](http://www.uniprot.org/citations/21477596), PubMed: [23949095](http://www.uniprot.org/citations/23949095)). Thereby, acts as a central switch between the signaling pathways activated by these second messengers with different cellular targets and opposite effects in numerous biological processes

(PubMed:8626589, PubMed:15544348). Also plays an important role in the biosynthesis of complex lipids (PubMed:8626589). Displays specificity for diacylglycerol substrates with an arachidonoyl acyl chain at the sn-2 position, with the highest activity toward 1-octadecanoyl-2- (5Z,8Z,11Z,14Z-eicosatetraenoyl)-sn-glycerol the main diacylglycerol intermediate within the phosphatidylinositol turnover cycle (PubMed:19744926, PubMed:22108654, PubMed:23274426). Can also phosphorylate diacylglycerol substrates with a linoleoyl acyl chain at the sn-2 position but much less efficiently (PubMed:22108654).

Cellular Location

Membrane; Single- pass membrane protein. Cytoplasm

Tissue Location

Expressed predominantly in testis. Expressed in endothelium, platelets and podocytes (at protein level)

DGKE Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

DGKE Antibody (C-term) Blocking Peptide - Images

DGKE Antibody (C-term) Blocking Peptide - Background

Diacylglycerol kinases are thought to be involved mainly in the regeneration of phosphatidylinositol (PI) from diacylglycerol in the PI-cycle during cell signal transduction. When expressed in mammalian cells, DGKE shows specificity for arachidonoyl-containing diacylglycerol. DGKE is expressed predominantly in testis.

DGKE Antibody (C-term) Blocking Peptide - References

Tang, W., et al., Gene 239(1):185-192 (1999).Hart, T.C., et al., Genomics 56(2):233-235 (1999).Tang, W., et al., J. Biol. Chem. 271(17):10237-10241 (1996).