

DUSP9 Antibody (C-term) Blocking Peptide

Synthetic peptide Catalog # BP8452b

Specification

DUSP9 Antibody (C-term) Blocking Peptide - Product Information

Primary Accession

Q99956

DUSP9 Antibody (C-term) Blocking Peptide - Additional Information

Gene ID 1852

Other Names

Dual specificity protein phosphatase 9, Mitogen-activated protein kinase phosphatase 4, MAP kinase phosphatase 4, MKP-4, DUSP9, MKP4

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP8452b was selected from the C-term region of human DUSP9. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

DUSP9 Antibody (C-term) Blocking Peptide - Protein Information

Name DUSP9

Synonyms MKP4

Function

Inactivates MAP kinases. Has a specificity for the ERK family.

Cellular Location

Cytoplasm.

DUSP9 Antibody (C-term) Blocking Peptide - Protocols



Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides

DUSP9 Antibody (C-term) Blocking Peptide - Images

DUSP9 Antibody (C-term) Blocking Peptide - Background

DUSP9 is a member of the dual specificity protein phosphatase subfamily. These phosphatases inactivate their target kinases by dephosphorylating both the phosphoserine/threonine and phosphotyrosine residues. They negatively regulate members of the mitogen-activated protein (MAP) kinase superfamily (MAPK/ERK, SAPK/JNK, p38), which is associated with cellular proliferation and differentiation. Different members of the family of dual specificity phosphatases show distinct substrate specificities for various MAP kinases, different tissue distribution and subcellular localization, and different modes of inducibility of their expression by extracellular stimuli. This DUSP9 shows selectivity for members of the ERK family of MAP kinases, is expressed only in placenta, kidney, and fetal liver, and is localized to the cytoplasm and nucleus.

DUSP9 Antibody (C-term) Blocking Peptide - References

Brenner, V., et al., Genomics 44(1):8-14 (1997). Muda, M., et al., J. Biol. Chem. 272(8):5141-5151 (1997).