

EIF3S2 Antibody (N-term) Blocking Peptide
Synthetic peptide
Catalog # BP8547a**Specification**

EIF3S2 Antibody (N-term) Blocking Peptide - Product InformationPrimary Accession [Q13347](#)**EIF3S2 Antibody (N-term) Blocking Peptide - Additional Information****Gene ID** 8668**Other Names**

Eukaryotic translation initiation factor 3 subunit I {ECO:0000255|HAMAP-Rule:MF_03008}, eIF3i {ECO:0000255|HAMAP-Rule:MF_03008}, Eukaryotic translation initiation factor 3 subunit 2 {ECO:0000255|HAMAP-Rule:MF_03008}, TGF-beta receptor-interacting protein 1, TRIP-1, eIF-3-beta {ECO:0000255|HAMAP-Rule:MF_03008}, eIF3 p36 {ECO:0000255|HAMAP-Rule:MF_03008}, EIF3I {ECO:0000255|HAMAP-Rule:MF_03008}

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP8547a](/products/AP8547a) was selected from the N-term region of human EIF3S2. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

EIF3S2 Antibody (N-term) Blocking Peptide - Protein Information**Name** EIF3I {ECO:0000255|HAMAP-Rule:MF_03008}**Function**

Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis (PubMed:[17581632](http://www.uniprot.org/citations/17581632), PubMed:[25849773](http://www.uniprot.org/citations/25849773), PubMed:[27462815](http://www.uniprot.org/citations/27462815)). The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl- tRNAi and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG

recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation (PubMed:17581632). The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression (PubMed:25849773).

Cellular Location

Cytoplasm {ECO:0000255|HAMAP-Rule:MF_03008}.

EIF3S2 Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

EIF3S2 Antibody (N-term) Blocking Peptide - Images**EIF3S2 Antibody (N-term) Blocking Peptide - Background**

EIF3S2 is the largest of the EIFs. It consists of at least 10 nonidentical subunits in mammals. In *S. cerevisiae* the p39 subunit contains WD repeats; these are thought to mediate protein-protein interactions. The p39 protein appears to be essential for maintaining the integrity of the yeast EIF3 complex. The mammalian EIF3-p36 subunit is homologous to yeast p39.

EIF3S2 Antibody (N-term) Blocking Peptide - References

Navarro,A.,et.al., Am. J. Physiol. Lung Cell Mol. Physiol. 296 (6), L928-L935 (2009)
Perard,J.,et.al.,FEBS Lett. 583 (1), 70-74 (2009)