

RPGRIP1L Antibody (N-term) Blocking Peptide
Synthetic peptide
Catalog # BP8567a**Specification**

RPGRIP1L Antibody (N-term) Blocking Peptide - Product InformationPrimary Accession [Q68CZ1](#)**RPGRIP1L Antibody (N-term) Blocking Peptide - Additional Information****Gene ID** 23322**Other Names**

Protein fantom, Nephrocystin-8, RPGR-interacting protein 1-like protein, RPGRIP1-like protein, RPGRIP1L, FTM, KIAA1005, NPHP8

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP8567a](/products/AP8567a) was selected from the N-term region of human RPGRIP1L. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

RPGRIP1L Antibody (N-term) Blocking Peptide - Protein Information**Name** RPGRIP1L**Synonyms** FTM, KIAA1005, NPHP8**Function**

Negatively regulates signaling through the G-protein coupled thromboxane A2 receptor (TBXA2R) (PubMed: [19464661](http://www.uniprot.org/citations/19464661)). May be involved in mechanisms like programmed cell death, craniofacial development, patterning of the limbs, and formation of the left-right axis (By similarity). Involved in the organization of apical junctions; the function is proposed to implicate a NPHP1-4-8 module. Does not seem to be strictly required for ciliogenesis (PubMed: [19464661](http://www.uniprot.org/citations/19464661)). Involved in establishment of planar cell polarity such as in cochlear sensory epithelium and is proposed to implicate stabilization of disheveled proteins (By similarity). Involved in regulation of proteasomal activity at the primary cilium probably implicating

association with PSDM2 (By similarity).

Cellular Location

Cytoplasm. Cytoplasm, cytoskeleton, cilium basal body {ECO:0000250|UniProtKB:Q8CG73, ECO:0000269|PubMed:21685204} Cytoplasm, cytoskeleton, cilium axoneme. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome {ECO:0000250|UniProtKB:Q8CG73}. Cell junction, tight junction {ECO:0000250|UniProtKB:Q8CG73}. Note=In cultured renal cells, it localizes diffusely in the cytoplasm but, as cells approach confluence, it accumulates to basolateral tight junctions. Localizes to the ciliary transition zone. {ECO:0000250|UniProtKB:Q8CG73}

Tissue Location

Ubiquitously expressed with relatively high level of expression in hypothalamus and islet. During early development, expressed in multiple organs including brain, eye, forelimb and kidney

RPGRIP1L Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

RPGRIP1L Antibody (N-term) Blocking Peptide - Images

RPGRIP1L Antibody (N-term) Blocking Peptide - Background

RPGRIP1L can localize to the basal body-centrosome complex or to primary cilia and centrosomes in ciliated cells. This protein has been found to interact with nephrocystin-4. Defects in this gene are a cause of Joubert syndrome type 7 (JBTS7) and Meckel syndrome type 5 (MKS5).

RPGRIP1L Antibody (N-term) Blocking Peptide - References

Brancati,F., et.al., Clin. Genet. 74 (2), 164-170 (2008)Stratigopoulos,G., et.al., Am. J. Physiol. Regul. Integr. Comp. Physiol. 294 (4), R1185-R1196(2008)