

# NQO2 Antibody (Center) Blocking Peptide

Synthetic peptide Catalog # BP8618c

# **Specification**

# NQO2 Antibody (Center) Blocking Peptide - Product Information

**Primary Accession** 

P16083

# NQO2 Antibody (Center) Blocking Peptide - Additional Information

**Gene ID 4835** 

#### **Other Names**

Ribosyldihydronicotinamide dehydrogenase [quinone], NRH dehydrogenase [quinone] 2, NRH:quinone oxidoreductase 2, Quinone reductase 2, QR2, NQO2, NMOR2

# **Target/Specificity**

The synthetic peptide sequence used to generate the antibody <a href=/products/AP8618c>AP8618c</a> was selected from the Center region of human NQO2. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

### NQO2 Antibody (Center) Blocking Peptide - Protein Information

Name NQO2

Synonyms NMOR2

### **Function**

The enzyme apparently serves as a quinone reductase in connection with conjugation reactions of hydroquinones involved in detoxification pathways as well as in biosynthetic processes such as the vitamin K-dependent gamma-carboxylation of glutamate residues in prothrombin synthesis.

### **Cellular Location**

Cytoplasm.



Tel: 858.875.1900 Fax: 858.875.1999

# NQO2 Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

## • Blocking Peptides

NQO2 Antibody (Center) Blocking Peptide - Images

## NQO2 Antibody (Center) Blocking Peptide - Background

NQO2 apparently serves as a quinone reductase in connection with conjugation reactions of hydroquinones involved in detoxification pathways as well as in biosynthetic processes such as the vitamin K-dependent gamma-carboxylation of glutamate residues in prothrombin synthesis.

# NQO2 Antibody (Center) Blocking Peptide - References

Foster, C.E., et.al., Biochemistry 38 (31), 9881-9886 (1999) Wu, K., Knox, R., et.al., Arch. Biochem. Biophys. 347 (2), 221-228 (1997)