10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

EHHADH Antibody (C-term) Blocking Peptide

Synthetic peptide
Catalog # BP8636b

Specification

EHHADH Antibody (C-term) Blocking Peptide - Product Information

Primary Accession Q08426

EHHADH Antibody (C-term) Blocking Peptide - Additional Information

Gene ID 1962

Other Names
Peroxisomal bifunctional enzyme, PBE, PBFE, Enoyl-CoA hydratase/3, 2-trans-enoyl-CoA isomerase, 3-hydroxyacyl-CoA dehydrogenase, EHHADH, ECHD

Target/Specificity
The synthetic peptide sequence used to generate the antibody AP8636b was selected from the C-term region of human EHHADH. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format
Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage
Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions
This product is for research use only. Not for use in diagnostic or therapeutic procedures.

EHHADH Antibody (C-term) Blocking Peptide - Protein Information

Name EHHADH (HGNC:3247)

Synonyms ECHD

Function
Peroxisomal trifunctional enzyme possessing 2-enoyl-CoA hydratase, 3-hydroxyacyl-CoA dehydrogenase, and delta 3, delta 2-enoyl- CoA isomerase activities. Catalyzes two of the four reactions of the long chain fatty acids peroxisomal beta-oxidation pathway (By similarity). Can also use branched-chain fatty acids such as 2-methyl- 2E-butenoyl-CoA as a substrate, which is hydrated into (2S,3S)-3- hydroxy-2-methylbutanoyl-CoA (By similarity). Optimal isomerase for 2,5 double bonds into 3,5 form isomerization in a range of enoyl-CoA species (Probable). Also able to isomerize both 3-cis and 3-trans double bonds into the 2-trans form in a range of enoyl-CoA species (By similarity). With HSD17B4, catalyzes the hydration of trans-2-enoyl-CoA and the dehydrogenation of 3-hydroxyacyl-CoA, but with opposite chiral specificity (PubMed:<a

href="http://www.uniprot.org/citations/15060085" target="_blank">15060085). Regulates the amount of medium-chain dicarboxylic fatty acids which are essential regulators of all fatty acid oxidation pathways (By similarity). Also involved in the degradation of long-chain dicarboxylic acids through peroxisomal beta- oxidation (PubMed:15060085).

Cellular Location
Peroxisome.

Tissue Location
Liver and kidney. Strongly expressed in the terminal segments of the proximal tubule. Lower amounts seen in the brain.

EHHADH Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Blocking Peptides

EHHADH Antibody (C-term) Blocking Peptide - Images

EHHADH Antibody (C-term) Blocking Peptide - Background

EHHADH is a bifunctional enzyme and is one of the four enzymes of the peroxisomal beta-oxidation pathway. The N-terminal region of the encoded protein contains enoyl-CoA hydratase activity while the C-terminal region contains 3-hydroxyacyl-CoA dehydrogenase activity.

EHHADH Antibody (C-term) Blocking Peptide - References

Chen,G.L., et.al., Biochem. Biophys. Res. Commun. 178 (3), 1084-1091 (1991)Lu,Y., et.al., J. Lipid Res. 49 (12), 2582-2589 (2008)