

COL18A1 Antibody (N-term) Blocking Peptide

Synthetic peptide Catalog # BP8852a

Specification

COL18A1 Antibody (N-term) Blocking Peptide - Product Information

Primary Accession

P39060

COL18A1 Antibody (N-term) Blocking Peptide - Additional Information

Gene ID 80781

Other Names

Collagen alpha-1(XVIII) chain, Endostatin, COL18A1

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP8852a was selected from the N-term region of human COL18A1. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

COL18A1 Antibody (N-term) Blocking Peptide - Protein Information

Name COL18A1 (HGNC:2195)

Function

Probably plays a major role in determining the retinal structure as well as in the closure of the neural tube.

Cellular Location

Secreted, extracellular space, extracellular matrix. Secreted, extracellular space, extracellular matrix, basement membrane {ECO:0000250|UniProtKB:P39061} [Endostatin]: Secreted. Secreted, extracellular space, extracellular matrix, basement membrane

Tissue Location

Detected in placenta (at protein level) (PubMed:32337544). Present in multiple organs with highest levels in liver, lung and kidney.



COL18A1 Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides

COL18A1 Antibody (N-term) Blocking Peptide - Images

COL18A1 Antibody (N-term) Blocking Peptide - Background

COL18A1 is encoding the alpha chain of type XVIII collagen. This collagen is one of the multiplexins, extracellular matrix proteins that contain multiple triple-helix domains (collagenous domains) interrupted by non-collagenous domains. The proteolytically produced C-terminal fragment of type XVIII collagen is endostatin, a potent antiangiogenic protein.

COL18A1 Antibody (N-term) Blocking Peptide - References

Fu, Y., et.al., Biochemistry 48 (49), 11655-11663 (2009)