

C1QB Antibody (N-term) Blocking Peptide
Synthetic peptide
Catalog # BP8893a**Specification**

C1QB Antibody (N-term) Blocking Peptide - Product InformationPrimary Accession [P02746](#)**C1QB Antibody (N-term) Blocking Peptide - Additional Information****Gene ID** 713**Other Names**

Complement C1q subcomponent subunit B, C1QB

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP8893a](/products/AP8893a) was selected from the N-term region of human C1QB. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

C1QB Antibody (N-term) Blocking Peptide - Protein Information**Name** C1QB**Function**

C1q associates with the proenzymes C1r and C1s to yield C1, the first component of the serum complement system. The collagen-like regions of C1q interact with the Ca(2+)-dependent C1r(2)C1s(2) proenzyme complex, and efficient activation of C1 takes place on interaction of the globular heads of C1q with the Fc regions of IgG or IgM antibody present in immune complexes.

Cellular Location

Secreted.

C1QB Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

C1QB Antibody (N-term) Blocking Peptide - Images

C1QB Antibody (N-term) Blocking Peptide - Background

C1QB is a major constituent of the human complement subcomponent C1q. C1q associates with C1r and C1s in order to yield the first component of the serum complement system. Deficiency of C1q has been associated with lupus erythematosus and glomerulonephritis. C1q is composed of 18 polypeptide chains: six A-chains, six B-chains, and six C-chains. Each chain contains a collagen-like region located near the N terminus and a C-terminal globular region. The A-, B-, and C-chains are arranged in the order A-C-B on chromosome 1.

C1QB Antibody (N-term) Blocking Peptide - References

Reid,K.B. et.al., Biochem. J. 173 (3), 863-868 (1978)Reid,K.B. et.al., Biochem. J. 179 (2), 367-371 (1979)